

INSTRUCTOR'S
SOLUTIONS MANUAL


DIACRITECH

INTERMEDIATE ALGEBRA:
FUNCTIONS AND AUTHENTIC
APPLICATIONS
SIXTH EDITION

Jay Lehmann

College of San Mateo





This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2019, 2015, 2011 Pearson Education, Inc.
Publishing as Pearson, 330 Hudson Street, NY NY 10013

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-478138-9

ISBN-10: 0-13-478138-4

Contents

Chapter 1 Linear Equations and Linear Functions

1.1	Using Qualitative Graphs to Describe Situations	1
1.2	Graphing Linear Equations	3
1.3	Slope of a Line	8
1.4	Meaning of Slope for Equations, Graphs, and Tables	12
1.5	Finding Linear Equations	17
1.6	Functions	23
	Chapter 1 Review Exercises	25
	Chapter 1 Test	29

Chapter 2 Modeling with Linear Functions

2.1	Using Lines to Model Data	33
2.2	Finding Equations of Linear Models	35
2.3	Function Notation and Making Predictions	39
2.4	Slope Is a Rate of Change	46
	Chapter 2 Review Exercises	51
	Chapter 2 Test	55

Chapter 3 Systems of Linear Equations and Systems of Linear Inequalities

3.1	Using Graphs and Tables to Solve Systems	58
3.2	Using Substitution and Elimination to Solve Systems	63
3.3	Using Systems to Model Data	70
3.4	Value, Interest, and Mixture Problems	75
3.5	Using Linear Inequalities in One Variable to Make Predictions	80
3.6	Linear Inequalities in Two Variables; Systems of Linear Inequalities	85
	Chapter 3 Review Exercises	90
	Chapter 3 Test	98
	Cumulative Review of Chapters 1–3	103

Chapter 4 Exponential Functions

4.1	Properties of Exponents	110
4.2	Rational Exponents	114
4.3	Graphing Exponential Functions	117
4.4	Finding Equations of Exponential Functions	122
4.5	Using Exponential Functions to Model Data	128
	Chapter 4 Review Exercises	135
	Chapter 4 Test	139

Chapter 5 Logarithmic Functions

5.1	Composite Functions	142
5.2	Inverse Functions	147
5.3	Logarithmic Functions	153
5.4	Properties of Logarithms	156
5.5	Using the Power Property with Exponential Models to Make Predictions	161
5.6	More Properties of Logarithms	168
5.7	Natural Logarithm	172
	Chapter 5 Review Exercises	177
	Chapter 5 Test	184
	Cumulative Review of Chapters 1–5	187

Chapter 6 Polynomial Functions

6.1	Adding and Subtracting Polynomial Expressions and Functions	194
6.2	Multiplying Polynomial Expressions and Functions	196
6.3	Dividing Polynomials: Long Division and Synthetic Division	201
6.4	Factoring Trinomials of the Form $x^2 + bx + c$; Factoring Out the GCF	206
6.5	Factoring Polynomials	208
6.6	Factoring Special Binomials; A Factoring Strategy	210
6.7	Using Factoring to Solve Polynomial Equations	212
	Chapter 6 Review Exercises	217
	Chapter 6 Test	220

Chapter 7 Quadratic Functions

7.1	Graphing Quadratic Functions in Vertex Form	223
7.2	Graphing Quadratic Functions in Standard Form	228
7.3	Using the Square Root Property to Solve Quadratic Equations	237
7.4	Solving Quadratic Equations by Completing the Square	243
7.5	Using the Quadratic Formula to Solve Quadratic Equations	249
7.6	Solving Systems of Linear Equations in Three Variables; Finding Quadratic Functions	259
7.7	Finding Quadratic Models	266
7.8	Modeling with Quadratic Functions	270
	Chapter 7 Review Exercises	275
	Chapter 7 Test	282
	Cumulative Review of Chapters 1–7	287

Chapter 8 Rational Functions

8.1	Finding the Domains of Rational Functions and Simplifying Rational Expressions	297
8.2	Multiplying and Dividing Rational Expressions; Converting Units	301
8.3	Adding and Subtracting Rational Expressions	307
8.4	Simplifying Complex Rational Expressions	312
8.5	Solving Rational Equations	317
8.6	Modeling with Rational Functions	326

8.7	Variation	331
	Chapter 8 Review Exercises	335
	Chapter 8 Test	343

Chapter 9 Radical Functions

9.1	Simplifying Radical Expressions	348
9.2	Adding, Subtracting, and Multiplying Radical Expressions	351
9.3	Rationalizing Denominators and Simplifying Quotients of Radical Expressions	355
9.4	Graphing and Combining Square Root Functions	360
9.5	Solving Radical Equations	365
9.6	Modeling with Square Root Functions	372
	Chapter 9 Review Exercises	378
	Chapter 9 Test	384

Chapter 10 Sequences and Series

10.1	Arithmetic Sequences	388
10.2	Geometric Sequences	390
10.3	Arithmetic Series	395
10.4	Geometric Series	398
	Chapter 10 Review Exercises	401
	Chapter 10 Test	403
	Cumulative Review of Chapters 1–10	405

Chapter 11 Additional Topics

11.1	Absolute Value: Equations and Inequalities	417
11.2	Performing Operations with Complex Numbers	421
11.3	Pythagorean Theorem, Distance Formula, and Circles	425
11.4	Ellipses and Hyperbolas	431
11.5	Solving Nonlinear Systems of Equations	440

Appendix A Reviewing Prerequisite Material

A.1	Plotting Points	448
A.2	Identifying Types of Numbers	448
A.3	Absolute Value	448
A.4	Performing Operations with Real Numbers	448
A.5	Exponents	448
A.6	Order of Operations	449
A.7	Constants, Variables, Expressions, and Equations	449
A.8	Distributive Law	449
A.9	Combining Like Terms	450
A.10	Solving Linear Equations in One Variable	450
A.11	Solving Equations in Two or More Variables	451
A.12	Equivalent Expressions and Equivalent Equations	452

Chapter 1

Linear Equations and Linear Functions

Homework 1.1

2. a. (c)
- b. (b)
- c. (a)
- d. (d)


4. The more pencils there are to buy, the more the total cost. Total cost c is the response variable and the number of pencils n is the explanatory variable.

6. The greater the rate water is added to a pool, the less time it takes to fill the pool. The number of hours to fill a pool t is the response variable and the rate water is added r is the explanatory variable.


8. The older the car in years, the greater the annual cost of repairs. The number of annual cost of repairs c is the response variable and the age of a car a is the explanatory variable.

10. The hotter the temperature, the more people go to the beach. The number of people at the beach n is the response variable and the temperature at the beach F is the explanatory variable.


12. The greater your annual income, the more federal taxes you pay. The federal taxes T is the response variable and the person's annual income I is the explanatory variable.



The response variable is s , the runner's speed, and the explanatory variable is t , time after she began her run. The graph shows her speed increases to a certain point and then decreases to a stop when $s = 0$. The rest of her time is spent walking at a moderate but slightly increasing speed.




Height h is the response variable and time, t is the explanatory variable. The height of the tennis ball decreases as it drops. The height then increases after the ball bounces from $h = 0$. This pattern is repeated with the maximum height decreasing after each bounce until the ball stops at $h = 0$.



P , the percentage of smokers, is the response variable and t , time in years since 2010, is the explanatory variable. This graph shows that the percentage of smokers approximately steadily decreases as time increases from 2010.




The foreign born percentage, p , is the response variable and time since 1910, t , is the explanatory variable. The graph shows that the percentage of the U.S. population that is foreign born decreased from 1910 to 1970 and then increased over time.




Volume V is the response variable and time t is the explanatory variable. The graph shows the volume of air in a person's lungs alternately increases and decreases as air is taken in and out over time.

2 ISM: Intermediate Algebra




Angle A is the response variable and minutes t is the explanatory variable. The graph shows that as the number of minutes past midnight increases, the angle (in degrees) between the hour hand and the minute hand of a clock increases.




The response variable is n (species in existence) and the explanatory variable is d (amount of deforestation). The graph shows that the number of species in existence decreases steadily due to increased deforestation.


28. Speed, s , is the response variable and current in miles per hour, c , is the explanatory variable.




The graph shows that a speedboat travels faster downstream as the river current increases.




The graph shows that a speedboat travels slower upstream as the “opposing” river current increases.




The percentage of times a person hits the bull’s-eye p is the response variable and the distance in feet from the dartboard d is the explanatory variable. The graph shows that as the distance in feet increases, the percentage of bull’s-eye hits decreases.



The number of hours to paint a house T is the response variable and the number of people on the painting crew N is the explanatory variable. The graph shows that as the number of people on the crew increases, the number of hours to paint decreases.



The driving time in hours T is the response variable and the speed in miles per hour the person drives S is the explanatory variable. The graph shows that as the speed in miles per hour increases, the driving time decreases.




Pressure P is the response variable and volume V is the explanatory variable. The graph shows that pressure in the balloon is greater when the volume is lower.

38. An increasing curve can have at most two intercepts where it crosses each axis.


40. First, determine what the response variable is and place it on the vertical axis. Then place the explanatory variable on the horizontal axis. There is no need for scaling on these axes. Determine from the given situation how to describe the relationship between the two variables. Use this relationship to see if an increasing or decreasing line or curve is necessary. Note where a point of intersection is needed on one or both axes.

Homework 1.2


2.




4.




6.




8.




10.



12.




14.



16. $0 = 4y - 20x$


$$\frac{20x}{4} = \frac{4y}{4}$$

$$5x = y$$




4 ISM: Intermediate Algebra


18. $10x - 5y = 20$
 $\frac{-5y}{-5} = \frac{-10x + 20}{-5}$
 $y = 2x - 4$




24. $6y - 4x - 1 = 7y - 2x - 4$
 $-y = 2x - 3$
 $\frac{-y}{-1} = \frac{2x - 3}{-1}$
 $y = -2x + 3$




20. $30x + 6y - 12 = 0$
 $\frac{6y}{6} = \frac{-30x + 12}{6}$
 $y = -5x + 2$




26. $2(y - 3) = 4(x + 1)$
 $2y - 6 = 4x + 4$
 $\frac{2y}{2} = \frac{4x + 10}{2}$
 $y = 2x + 5$




22. $3y + 3x - 2 = 7$
 $\frac{3y}{3} = \frac{-3x + 9}{3}$
 $y = -x + 3$




28. $5x - 2(3y - 1) = -2y - 3(x - 2)$
 $5x - 6y + 2 = -2y - 3x + 6$
 $-6y + 2y = -3x - 5x + 4$
 $\frac{-4y}{-4} = \frac{-8x + 4}{-4}$
 $y = 2x - 1$




30.



iii.




32.




b. An equation of the form $x = a$ is a vertical line passing through $(a, 0)$.


38.




34.




40.




36. a. i.




42.



ii.



44. $2x - 3y = 6$
 $-3y = -2x + 6$
 $y = \frac{(-2x + 6)}{-3} = \frac{2}{3}x - 2$



6 ISM: Intermediate Algebra


$(0, -2)$ is a point on the line.

$$2x - 3y = 6$$

$$2(0) - 3(-2) \stackrel{?}{=} 6$$

$$0 + 6 \stackrel{?}{=} 6$$

$$6 \stackrel{?}{=} 6 \text{ true}$$



b. Answers may vary. Example:

x	y
0	1.890
0.2	1.404
0.4	0.918
0.6	0.432
0.8	-0.054

48. $3 - 5x - 2 = 4x + 9 - 7x$

$$-5x + 1 = -3x + 9$$

$$\frac{-2x}{-2} = \frac{8}{-2}$$

$$x = -4$$

50. $4(2t - 1) = -5t + 7$

$$8t - 4 = -5t + 7$$

$$\frac{13t}{13} = \frac{11}{13}$$

$$t = \frac{11}{13}$$

52. $7 - 2(5 - 4x) = 3x - (8 - 3x)$

$$7 - 10 + 8x = 3x - 8 + 3x$$

$$-3 + 8x = 6x - 8$$

$$\frac{2x}{2} = \frac{-5}{2}$$

$$x = -\frac{5}{2}$$

54. $-4(a + 6) + 5(a - 3) = 3(a - 1)$

$$-4a - 24 + 5a - 15 = 3a - 3$$

$$a - 39 = 3a - 3$$

$$\frac{-2a}{-2} = \frac{36}{-2}$$

$$a = -18$$

56. $\frac{1}{3}x - \frac{1}{4} = \frac{2}{3}$

$$12\left(\frac{1}{3}x - \frac{1}{4}\right) = 12\left(\frac{2}{3}\right)$$

$$4x - 3 = 8$$

$$\frac{4x}{4} = \frac{11}{4}$$

$$x = \frac{11}{4}$$

58. $-\frac{3}{4}w - \frac{5}{8} = \frac{3}{2}w + \frac{1}{4}$

$$8\left(-\frac{3}{4}w - \frac{5}{8}\right) = 8\left(\frac{3}{2}w + \frac{1}{4}\right)$$

$$-6w - 5 = 12w + 2$$

$$\frac{-18w}{-18} = \frac{7}{-18}$$

$$w = -\frac{7}{18}$$

60. $-6.54x + 87.35 = -4.66x - 99.03$

$$\frac{-1.88x}{-1.88} = \frac{-186.38}{-1.88}$$

$$x \approx 99.14$$

62. $ax + by = c$

$$\frac{by}{b} = \frac{c - ax}{b}$$

$$y = \frac{c - ax}{b}$$

64. $\frac{x}{a} + \frac{y}{a} = 1$

$$a\left(\frac{x}{a} + \frac{y}{a}\right) = a(1)$$

$$x + y = a$$

$$y = a - x$$

66. $y = -3x - 12$

To find the x -intercept, let $y = 0$ and solve for x .

$$0 = -3x - 12$$

$$12 = -3x$$

$$-4 = x$$

The x -intercept is $(-4, 0)$. To find the y -intercept, let $x = 0$ and solve for y .

$$y = -3(0) - 12 = 0 - 12 = -12$$

The y -intercept is $(0, -12)$.

68. $5x - 4y = 20$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} 5x - 4(0) &= 20 \\ 5x - 0 &= 20 \\ 5x &= 20 \\ x &= 4 \end{aligned}$$

The x -intercept is $(4, 0)$. To find the y -intercept, let $x = 0$ and solve for y .

$$\begin{aligned} 5(0) - 4y &= 20 \\ 0 - 4y &= 20 \\ -4y &= 20 \\ y &= -5 \end{aligned}$$

The y -intercept is $(0, -5)$.

70. $y = -2x$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} 0 &= -2x \\ 0 &= x \end{aligned}$$

The x -intercept is $(0, 0)$. This is also the y -intercept.

72. $x = -2$

Since $x = -2$ is a vertical line, it never intersects the y -axis. Therefore, there is no y -intercept.

Since the graph passes through $(-2, 0)$, this is the x -intercept.

74. $ax + by = c$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} ax + b(0) &= c \\ ax &= c \\ x &= \frac{c}{a} \end{aligned}$$

The x -intercept is $\left(\frac{c}{a}, 0\right)$.

To find the y -intercept, let $x = 0$ and solve for y .

$$\begin{aligned} a(0) + by &= c \\ by &= c \\ y &= \frac{c}{b} \end{aligned}$$

The y -intercept is $\left(0, \frac{c}{b}\right)$.

76. $a(x - by) = c$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} a(x - b \cdot 0) &= c \\ ax &= c \\ x &= \frac{c}{a} \end{aligned}$$

The x -intercept is $\left(\frac{c}{a}, 0\right)$.

To find the y -intercept, let $x = 0$ and solve for y .

$$\begin{aligned} a(0 - by) &= c \\ -aby &= c \\ y &= -\frac{c}{ab} \end{aligned}$$

The y -intercept is $\left(0, -\frac{c}{ab}\right)$.

78. $ay = b(c + dx)$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} a(0) &= b(c + dx) \\ 0 &= bc + bdx \\ -bdx &= bc \\ x &= -\frac{c}{d} \end{aligned}$$

The x -intercept is $\left(-\frac{c}{d}, 0\right)$.

To find the y -intercept, let $x = 0$ and solve for y .

$$\begin{aligned} ay &= b(c + d \cdot 0) \\ ay &= bc \\ y &= \frac{bc}{a} \end{aligned}$$

The y -intercept is $\left(0, \frac{bc}{a}\right)$.

80. $\frac{y - b}{m} = x$

To find the x -intercept, let $y = 0$ and solve for x .

$$\begin{aligned} \frac{(0) - b}{m} &= x \\ \frac{-b}{m} &= x \end{aligned}$$

The x -intercept is $\left(\frac{-b}{m}, 0\right)$.

To find the y -intercept, let $x = 0$ and solve for y .

$$\begin{aligned} \frac{y - b}{m} &= 0 \\ y - b &= 0 \\ y &= b \end{aligned}$$

The y -intercept is $(0, b)$.

82. Answers may vary. Example:

x	y
-3	-3
0	-1
1.5	0
3	1
6	3

84. $x = 2$

8 ISM: Intermediate Algebra

86. $y = 1$


88. $y \approx -\frac{1}{2}$

90. $x = 8$

92. $x = -6$

94. $x \approx -1$

96. a.



b. To find the y -intercept, let $x = 0$ and solve for y .

$$y = -2(0) + 10 = 0 + 10 = 10$$

The y -intercept is $(0, 10)$. This means that there are 10 gallons of gas in the tank at the time of refueling.

c. To find the x -intercept, let $y = 0$ and solve for x .

$$0 = -2x + 10$$

$$2x = 10$$

$$x = 5$$

The x -intercept is $(5, 0)$. This means that when 5 hours have passed since filling the tank, the tank will be empty.

98. The graph of $y + x^2 = 5x + x^2 + 1$ is a line. When combining like terms, note that the equation becomes $y = 5x + 1$. The x^2 term cancels out. It is now an equation in the form $y = mx + b$.

100. Since $(2, 11)$ is a point on the graph of the equation $y = mx + 3$, $(2, 11)$ satisfies the equation. Substitute $x = 2$ and $y = 11$ into $y = mx + 3$ and solve for m .

$$11 = m(2) + 3$$

$$8 = 2m$$

$$4 = m$$

Therefore, $m = 4$ in $y = mx + 3$.

102. a. Points A and F satisfy the equation $y = ax + b$ since these points lie on the line for this equation.

b. Points D and F satisfy the equation $y = cx + d$ since these points lie on the line for this equation.

c. Point F satisfies both equations since it lies on both lines. It lies at the point where the lines intersect.

d. Points B, C, and E do not satisfy either equation since they do not lie on either line.

104. The y -coordinate of the x -intercept is 0 because the x -intercept is a point on the x -axis. All points on the x -axis have a y -coordinate equal to 0. The x -coordinate of the y -intercept is 0 because the y -intercept is a point on the y -axis. All points on the y -axis have an x -coordinate equal to 0.

106. No, every line does not have an x -intercept. Answers may vary. Example: Most horizontal lines do not have x -intercepts. For example, the line $y = 1$ does not have an x -intercept.

Homework 1.3

2. $m_A = \frac{2500}{8000} = 0.3125$

$$m_B = \frac{3100}{9500} \approx 0.326$$

Airplane B is making a steeper climb than airplane A.

4. The slope of the ski run from the top of the mountain to the chairlift is $\frac{|-100|}{400} = \frac{1}{4}$. If the vertical distance from the top of the mountain to the chairlift is 100 yards and the vertical distance for the entire run is 415 yards, then the vertical distance from the chairlift to the restaurant is $415 - 100 = 315$ yards. Also, since the horizontal distance from the top of the mountain to the chairlift is 400 yards and the horizontal distance for the entire run is 1300 yards, the horizontal distance from the chairlift to the restaurant is $1300 - 400 = 900$ yards. Therefore, the slope of the ski run from the chairlift to the restaurant is

$$\frac{|-315|}{900} = \frac{315}{900} \text{ or } \frac{7}{20}$$

$$6. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - 4}{1 - 5} = \frac{4}{-4} = -1$$

Since m is negative 1, the line is decreasing.

$$8. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - (-2)}{2 - 8} = \frac{8}{-6} = -\frac{4}{3}$$

Since m is negative $\frac{4}{3}$, the line is decreasing.

$$10. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{6 - (-10)}{-3 - 1} = \frac{16}{-4} = -4$$

Since m is negative 4, the line is decreasing.

$$12. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-3 - (-9)}{2 - 6} = \frac{6}{-4} = -\frac{3}{2}$$

Since m is negative $\frac{3}{2}$, the line is decreasing.

$$14. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-12 - (-2)}{-1 - 7} = \frac{-10}{-8} = \frac{5}{4}$$

Since m is positive $\frac{5}{4}$, the line is increasing.

$$16. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-6 - (-5)}{-1 - (-2)} = \frac{-6 + 5}{-1 + 2} = \frac{-1}{1} = -1$$

Since m is negative 1, the line is decreasing.

$$18. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{100 - 0}{100 - 0} = \frac{100}{100} = 1$$

Since m is positive 1, the line is increasing.

$$20. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - (-1)}{-3 - 5} = \frac{0}{-8} = 0$$

Since m is zero, the line is horizontal.

$$22. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 8}{4 - 4} = \frac{-9}{0} = \text{undefined}$$

Since m is undefined, the line is vertical.

$$24. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{0 - (-3)}{-6 - 0} = \frac{3}{-6} = -\frac{1}{2}$$

Since m is negative $\frac{1}{2}$, the line is decreasing.

$$26. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-7.6 - 2.2}{-5.1 - (-3.9)} = \frac{-9.8}{-1.2} = 8.17$$

Since m is positive 8.17, the line is increasing.

$$28. m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{82.78 - (-66.66)}{-25.41 - (-11.26)} = \frac{149.44}{-14.15} = -10.56$$

Since m is negative 10.56, the line is decreasing.

30. The points (0, 4) and (-2, 1) lie on the line.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{4 - 1}{0 - (-2)} = \frac{3}{2}$$

The slope of the line is $\frac{3}{2}$.

32. Since $m_2 = m_1$, the lines are parallel.

34. Since $m_2 = -\frac{1}{m_1}$, the lines are perpendicular.


36. Since $m_2 = -\frac{1}{m_1}$, the lines are perpendicular.

38. Since $m_2 = m_1$, the lines are parallel.


40. Since m_1 and m_2 are undefined, l_1 and l_2 are vertical lines. Vertical lines are parallel.

42. The two lines are not parallel, since their slopes, $\frac{1}{3}$ and $\frac{3}{10}$, are not equal.

44. Answers may vary. Example:




46. Answers may vary. Example:




10 ISM: Intermediate Algebra


48. Answers may vary. Example:



50. Answers may vary. Example:




52.



Answers may vary. Example:


Since $\frac{-2}{6} = -\frac{1}{3}$ and $\frac{1}{-3} = -\frac{1}{3}$, the slopes are the same. Both lines have the same slope and the same y -intercept, so they must be the same line.

54. Answers may vary. Example:



The lines have the same steepness.


56. Answers may vary. Example:



The points $(0, 0)$ and $(10, -1)$ are on the line and can be used to find the slope.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 0}{10 - 0} = \frac{-1}{10} = -\frac{1}{10}$$


58. Answers may vary. Example:



The points $(0, 0)$ and $(1, 10)$ are on the line and can be used to find the slope.


$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{10 - 0}{1 - 0} = \frac{10}{1} = 10$$

60. Answers may vary. Example:



The line must be increasing and almost horizontal.

62. Answers may vary. Example:




The line must be decreasing and almost vertical.

64. The student exchanged y_1 with y_2 in calculating the slope. The correct slope is negative instead of positive since

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 9}{7 - 2} = \frac{-8}{5} = -\frac{8}{5}.$$

66.



Answers may vary. Example: Three points that lie on the line are (3, 11), (0, 6), and (-9, -9) since the line equation is

$$y = \frac{5}{3}x + 6.$$

68. a. $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{5 - 3}{7 - 2} = \frac{2}{5}$


b. $m = \frac{y_1 - y_2}{x_1 - x_2} = \frac{3 - 5}{2 - 7} = \frac{-2}{-5} = \frac{2}{5}$

c. The results are the same. It does not matter which point is taken first.

d. $\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_1 - y_2}{x_1 - x_2}$
 $\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_2 - y_1}{x_2 - x_1} \cdot \frac{-1}{-1}$
 $\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_2 - y_1}{x_2 - x_1} \cdot 1$
 $\frac{y_2 - y_1}{x_2 - x_1} = \frac{y_2 - y_1}{x_2 - x_1}$


e. It does not matter which point is used first and which point is used second. The slope will be the same regardless of which point is considered first.

70. a.




There are two possibilities for the other two vertices, since the figure is a square and sides must have equal length. The coordinates for the vertices are (-3, 1) and (-3, 7). Another possibility is (9, 1) and (9, 7).

b. One possibility is the point (0, -3).




A second possibility is the point (6, 5).



Another possibility is (-14, -1).

72. Answers may vary. Example:




74. Answers may vary.


12 ISM: Intermediate Algebra

Homework 1.4


2. Since $y = 3x + 6$ is in the form $y = mx + b$, the slope is $m = 3 = \frac{3}{1} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, 6)$.




4. Since $y = -3x + 5$ is in the form $y = mx + b$, the slope is $m = -3 = \frac{-3}{1} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, 5)$.




6. Since $y = \frac{1}{2}x - 8$ is in the form $y = mx + b$, the slope is $m = \frac{1}{2} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, -8)$.



8. Since $y = -\frac{4}{3}x + 1$ is in the form $y = mx + b$, the slope is $m = -\frac{4}{3} = \frac{-4}{3} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, 1)$.




10. Since $y = -\frac{2}{5}x - 3$ is in the form $y = mx + b$, the slope is $m = -\frac{2}{5} = \frac{-2}{5} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, -3)$.



12. First, rewrite $2x + y = -1$ in slope-intercept form.

$$\begin{aligned} 2x + y &= -1 \\ y &= -2x - 1 \end{aligned}$$


- The slope is $m = -2 = \frac{-2}{1} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, -1)$.



14. First, rewrite $3y - 5x = 6$ in slope-intercept form.

$$\begin{aligned} 3y - 5x &= 6 \\ 3y &= 5x + 6 \\ y &= \frac{5}{3}x + 2 \end{aligned}$$


The slope is $m = \frac{5}{3} = \frac{\text{rise}}{\text{run}}$ and the y -intercept is $(0, 2)$.



16. First, rewrite $2(y - 3x) = 8$ in slope-intercept form.

$$\begin{aligned} 2(y - 3x) &= 8 \\ 2y - 6x &= 8 \\ 2y &= 8 + 6x \\ y &= 4 + 3x \end{aligned}$$


The slope is $m = 3 = \frac{3}{1} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, 4)$.



18. First, rewrite $-5x - 15y + 23 = 3$ in slope-intercept form.

$$\begin{aligned} -5x - 15y + 23 &= 3 \\ -5x - 15y &= -20 \\ -15y &= 5x - 20 \\ y &= -\frac{1}{3}x + \frac{4}{3} \end{aligned}$$


The slope is $m = -\frac{1}{3} = \frac{-1}{3} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, \frac{4}{3})$.



20. First, rewrite $3y - 6x + 2 = 7y - x - 6$ in slope-intercept form.

$$\begin{aligned} 3y - 6x + 2 &= 7y - x - 6 \\ -4y &= 5x - 8 \\ y &= -\frac{5}{4}x + 2 \end{aligned}$$


The slope is $m = -\frac{5}{4} = \frac{-5}{4} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, 2)$.



22. First, rewrite $8 - 2(y - 3x) = 2 + 4(x - 2y)$ in slope-intercept form.

$$\begin{aligned} 8 - 2(y - 3x) &= 2 + 4(x - 2y) \\ 8 - 2y + 6x &= 2 + 4x - 8y \\ 6y &= -2x - 6 \\ y &= -\frac{1}{3}x - 1 \end{aligned}$$


The slope is $m = -\frac{1}{3} = \frac{-1}{3} = \frac{\text{rise}}{\text{run}}$, and the y -intercept is $(0, -1)$.



14 ISM: Intermediate Algebra

24. Rewrite $y = -7x$ as $y = -7x + 0$ to obtain slope-intercept form. The slope is


$$m = -7 = \frac{-7}{1} = \frac{\text{rise}}{\text{run}}, \text{ and the } y\text{-intercept is } (0, 0).$$



26. Since $y = 0.25x - 2$ is in the form


$$y = mx + b, \text{ the slope is } m = 0.25 = \frac{1}{4} = \frac{\text{rise}}{\text{run}},$$

and the y -intercept is $(0, -2)$.




28. Rewrite $y = -x$ as $y = -1x + 0$ to obtain slope-intercept form. The slope is

$$m = -1 = \frac{-1}{1} = \frac{\text{rise}}{\text{run}}, \text{ and the } y\text{-intercept is } (0, 0).$$



30. The linear equation $y = 0$ is a horizontal line. The slope of a horizontal line is $m = 0$. Since this is the equation of the x -axis, there are an infinite number of x -intercepts. The y -intercept is at $(0, 0)$.




32. Solve for y .

$$y - 3 = 0$$

$$y = 3$$

The linear equation $y = 3$ is a horizontal line.

The slope of a horizontal line is $m = 0$, and there is no x -intercept. The y -intercept is $(0, 3)$.



34. Solve for y .

$$ax + by + c = 0$$

$$by = -ax - c$$

$$y = -\frac{a}{b}x - \frac{c}{b}$$

The slope is $-\frac{a}{b}$ and the y -intercept is

$$\left(0, -\frac{c}{b}\right).$$

36. Solve for y .

$$ay = b(x + d)$$

$$ay = bx + bd$$

$$y = \frac{b}{a}x + \frac{bd}{a}$$

The slope is $\frac{b}{a}$ and the y -intercept is

$$\left(0, \frac{bd}{a}\right).$$

38. Solve for y .

$$a(y - b) = x$$

$$y - b = \frac{1}{a}x$$

$$y = \frac{1}{a}x + b$$

The slope is $\frac{1}{a}$ and the y -intercept is $(0, b)$.

40. Solve for y .

$$a(x + y) = d$$

$$x + y = \frac{d}{a}$$

$$y = -x + \frac{d}{a}$$

The slope is -1 and the y -intercept is $(0, \frac{d}{a})$.

42. Solve for y .

$$\frac{y + b}{a} = x$$

$$y + b = ax$$

$$y = ax - b$$

The slope is a and the y -intercept is $(0, -b)$.

44. Set 1 is linear. As x increases by 1, y

decreases by 3. The slope is $\frac{-3}{1} = -3$.

Set 2 is linear. As x increases by 2, y

increases by 3. The slope is $\frac{3}{2}$.

Set 3 is linear. Regardless of the value of x , y is 8. All points lie on the line $y = 8$. The slope of the (horizontal) line is 0.

Set 4 is linear. Regardless of the value of y , x is 5. The points lie on the line $x = 5$. The slope of the (vertical) line is undefined.

	Eq. 1		Eq. 2		Eq. 3		Eq. 4	
	x	y	x	y	x	y	x	y
	0	16	0	-6	1	36	10	80
	1	23	1	-2	2	32	11	77.5
	2	30	2	2	3	28	12	75
	3	37	3	6	4	24	13	72.5
	4	44	4	10	5	20	14	70
	5	51	5	14	6	16	15	67.5

48. Since the slopes of 7 and -7 are neither the same nor negative reciprocals, the lines are neither parallel nor perpendicular.

50. Since $-\frac{5}{2}$ and $\frac{2}{5}$ are negative reciprocals, the lines are perpendicular.

52. Solve each equation for y .

$$4x + y = 6 \qquad x - 4y = 5$$

$$y = -4x + 6 \qquad -4y = -x + 5$$

$$y = \frac{1}{4}x - \frac{5}{4}$$

Since -4 and $\frac{1}{4}$ are negative reciprocals, the lines are perpendicular.

54. Solve each equation for y .

$$8x - 4y = 1 \qquad x + 2y = 6$$

$$-4y = -8x + 1 \qquad 2y = -x + 6$$

$$y = 2x - \frac{1}{4} \qquad y = -\frac{1}{2}x + 3$$

Since 2 and $-\frac{1}{2}$ are negative reciprocals, the lines are perpendicular.

56. $y = 5$ and $y = -2$ are both horizontal lines. Horizontal lines do not intersect. Therefore, these lines are parallel.

58. $x = -2$ is a vertical line and $y = -4$ is a horizontal line. All vertical and horizontal lines are perpendicular.

60. a. Substitute values for x in the equation $y = -400x + 2400$ to solve for y .
 When $x = 0$, $y = -400(0) + 2400 = 2400$.
 When $x = 1$, $y = -400(1) + 2400 = 2000$.
 Similar calculations yield the following table.

Time (minutes)	Altitude (feet)
x	y
0	2400
1	2000
2	1600
3	1200
4	800
5	400
6	0

- b. Each minute, the height of the balloon decreases by 400 feet. The slope of -400 in the equation shows this decrease of 400. As the value of the explanatory variable (time) increases by 1, the value of the

16 ISM: Intermediate Algebra

response variable (altitude) decreases by 400 as determined by the slope. After 6 minutes, the balloon will have descended to the ground.

62. a. Substitute values for x in the equation $y = 0.5x + 6$ to solve for y and complete the table.

When $x = 0$, $y = 0.5(0) + 6 = 6$.

When $x = 1$, $y = 0.5(1) + 6 = 0.5 + 6 = 6.5$.


Similar calculations yield the following table.

Number of Years College Has Been Open	Enrollment (thousands of students)
x	y
0	6.0
1	6.5
2	7.0
3	7.5
4	8.0

- b. Each year the college's enrollment increases by 500 students. This corresponds to the slope of $y = 0.5x + 6$, since $0.5(1000) = 500$. As the value of the explanatory variable (number of years that the college has been open) increases by 1, the value of the response variable (enrollment in thousands of students) increases by 500 as determined by the slope.

64. Answers may vary. Example:

$$\begin{array}{ll} y = x + 1 & y = x + 2 \\ y = x + 3 & y = x + 4 \\ y = x + 5 & \end{array}$$



66. The slope is $m = \frac{\text{rise}}{\text{run}} = \frac{2}{3}$. The y -intercept is $(0, -1)$. The equation of the line is

$$y = \frac{2}{3}x - 1.$$

68. Answers may vary. Example:
 $y = 500x - 3$. The slope is a large positive number, and the y -intercept is negative.

70. Answers may vary. Example:
 $y = -\frac{1}{200}x + 5$. The slope is a small negative number, and the y -intercept is positive.

72. a.

```
WINDOW
Xmin=-.047
Xmax=.047
Xscl=1
Ymin=-.031
Ymax=.031
Yscl=1
Xres=1
```

b.

```
WINDOW
Xmin=-10
Xmax=10
Xscl=1
Ymin=19900
Ymax=20100
Yscl=10
Xres=1
```

c.


```
WINDOW
Xmin=-10
Xmax=10
Xscl=1
Ymin=-10000
Ymax=10000
Yscl=1000
Xres=1
```

74. Since we have the slope, $m = -5$, and a point on the line, $(0, 4)$, we can substitute in the equation $y = mx + b$ to find the y -intercept, b .

$$\begin{aligned} y &= mx + b \\ 4 &= -5(0) + b \\ 4 &= b \end{aligned}$$


We can now write the equation of the line in slope-intercept form, $y = -5x + 4$.

76. a.




- b. From the graph in part (a), we see that the y -intercept, b , is -5 . Using the slope and the y -intercept, we can write the equation of the line in slope-intercept form.

$$\begin{aligned} y &= -6x + (-5) \\ y &= -6x - 5 \end{aligned}$$



78. a.



$$y = mx + b$$

$$y = \frac{1}{4}x + b$$

Substitute $(x, y) = (-3, 2)$


$$y = mx + b$$

$$2 = \frac{-3}{4} + b$$

$$b = \frac{11}{4}$$

- b. From part (a), we see that the y -intercept, b , is $2\frac{3}{4}$ or $\frac{11}{4}$. Using the slope and the y -intercept, we can write the equation of the line in slope–intercept form.

$$y = \frac{1}{4}x + \frac{11}{4}$$



80. a. Each line is a vertical line with an undefined slope.

- b. The slope of the graph for any linear equation of the form $x = k$ is undefined when k is a constant.

82. No. When the linear equation is written in slope–intercept form, the coefficient of x is the slope of a line. So, instead of $3x$, the slope is 3.

84. Answers may vary. Example:
 $y = 2x - 3$

$$\text{Let } x_1 = 1 : y_1 = 2(1) - 3 = 2 - 3 = -1$$

$$\text{Let } x_2 = -1 : y_2 = 2(-1) - 3 = -2 - 3 = -5$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-5 - (-1)}{-1 - 1} = \frac{-4}{-2} = 2$$

The slope is the same as m in the equation.

86. Answers may vary.

Homework 1.5

2. We are given the slope, $m = 2$, and a point on the line, $(3, 1)$. Use $y = mx + b$ to find b .

$$y = mx + b$$


$$1 = 2(3) + b$$

$$1 = 6 + b$$

$$-5 = b$$

Now, substitute for m and b in slope–intercept form to obtain the equation of the line.

$$y = 2x - 5$$



The sign of m (positive) agrees with the increasing line from the graphing calculator.

4. We are given the slope, $m = -4$, and a point on the line, $(-2, -8)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$-8 = -4(-2) + b$$


$$-8 = 8 + b$$

$$-16 = b$$

Now, substitute for m and b in slope–intercept form to obtain the equation of the line.

$$y = -4x + (-16)$$

$$y = -4x - 16$$



The sign of m (negative) agrees with the decreasing line from the graphing calculator.

6. We are given the slope, $m = \frac{2}{3}$, and a point on the line, $(6, 1)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$1 = \frac{2}{3}(6) + b$$


$$1 = 4 + b$$

$$-3 = b$$

Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$y = \frac{2}{3}x + (-3)$$

$$y = \frac{2}{3}x - 3$$



The sign of m (positive) agrees with the increasing line shown on the graphing calculator screen.

8. We are given the slope, $m = -\frac{1}{4}$, and a point

on the line, $(-2, 5)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$5 = -\frac{1}{4}(-2) + b$$


$$5 = \frac{1}{2} + b$$

$$\frac{10}{2} = \frac{1}{2} + b$$

$$\frac{9}{2} = b$$

Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$y = -\frac{1}{4}x + \frac{9}{2}$$



The sign of m (negative) agrees with the decreasing line shown on the graphing calculator screen.

10. We are given the slope, $m = -\frac{3}{4}$, and a point on the line, $(-5, -2)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$-2 = -\frac{3}{4}(-5) + b$$


$$-2 = \frac{15}{4} + b$$

$$-\frac{8}{4} = \frac{15}{4} + b$$

$$-\frac{23}{4} = b$$


Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$y = -\frac{3}{4}x - \frac{23}{4}$$




The sign of m (negative) agrees with the decreasing line shown on the graphing calculator screen.

12. We are given the slope, $m = 0$, and a point on the line, $(-3, -4)$. We know that a line with a slope of 0 is a horizontal line. Horizontal lines are of the form $y = b$. In this example, $y = -4$.



Since m is zero, the line is horizontal. This is shown on the graphing calculator screen.

14. We are given that m is undefined, and a point on the line is $(-5, 1)$. We know that a line with an undefined slope is a vertical line. Vertical lines are of the form $x = a$. In this example, $x = -5$.



Since m is undefined, the line is vertical. This is shown on the graphing calculator screen.

16. We are given the slope, $m = -2.7$, and a point on the line, $(6.2, -4.9)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$-4.9 = -2.7(6.2) + b$$

$$-4.9 = -16.74 + b$$

$$11.84 = b$$

Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$y = -2.7x + 11.84$$

To check, substitute $(6.2, -4.9)$ into the equation.

$$y = -2.7x + 11.84$$

$$-4.9 \stackrel{?}{=} -2.7(6.2) + 11.84$$

$$-4.9 \stackrel{?}{=} -16.74 + 11.84$$

$$-4.9 \stackrel{?}{=} -4.9 \text{ true}$$

18. We are given the slope, $m = 1.94$, and a point on the line, $(-2.53, -3.77)$. Use $y = mx + b$ to find b .

$$y = mx + b$$

$$-3.77 = 1.94(-2.53) + b$$

$$-3.77 = -4.9082 + b$$

$$1.14 \approx b$$

Substitute m and b in slope-intercept form to obtain the equation of the line.

$$y = 1.94x + 1.14$$

To check, substitute $(-2.53, -3.77)$ into the equation.

$$y = 1.94x + 1.14$$

$$-3.77 \stackrel{?}{=} 1.94(-2.53) + 1.14$$

$$-3.77 \stackrel{?}{=} -4.9082 + 1.14$$

$$-3.77 \stackrel{?}{=} -3.77 \text{ true}$$

20. First, find the slope.

$$m = \frac{5-1}{3-7} = \frac{4}{-4} = -1$$

So, $y = -x + b$. Since the line contains $(3, 5)$, substitute 3 for x and 5 for y and solve for b .

$$5 = -1(3) + b$$

$$5 = -3 + b$$

$$8 = b$$

So, the equation is $y = -x + 8$.

22. First, find the slope.

$$m = \frac{3-(-9)}{-1-2} = \frac{12}{-3} = -4$$

So, $y = -4x + b$. Since the line contains $(-1, 3)$, substitute -1 for x and 3 for y and solve for b .

$$3 = -4(-1) + b$$

$$3 = 4 + b$$

$$-1 = b$$

So, the equation is $y = -4x - 1$.

24. First, find the slope.

$$m = \frac{-1-(-7)}{-4-(-2)} = \frac{-1+7}{-4+2} = \frac{6}{-2} = -3$$

So, $y = -3x + b$. Since the line contains $(-4, -1)$, substitute -4 and for x and -1 for y solve for b .

$$-1 = -3(-4) + b$$

$$-1 = 12 + b$$

$$-13 = b$$

So, the equation is $y = -3x - 13$.

26. First, find the slope.

$$m = \frac{8-5}{0-4} = \frac{3}{-4} = -\frac{3}{4}$$

So, $y = -\frac{3}{4}x + b$. Since the line contains $(0, 8)$,

substitute 0 for x and 8 for y to solve for b .

$$8 = -\frac{3}{4}(0) + b$$

$$8 = b$$

So, the equation is $y = -\frac{3}{4}x + 8$.

28. First, find the slope.

$$m = \frac{9-2}{5-3} = \frac{7}{2}$$

So, $y = \frac{7}{2}x + b$. Since the line passes through

$(3, 2)$, substitute 3 for x and 2 for y to solve for b .

$$2 = \frac{7}{2}(3) + b$$

$$2 = \frac{21}{2} + b$$

$$\frac{4}{2} = \frac{21}{2} + b$$

$$-\frac{17}{2} = b$$

So, the equation is $y = \frac{7}{2}x - \frac{17}{2}$.

30. First, find the slope.

$$m = \frac{-1-(-3)}{2-5} = \frac{2}{-3} = -\frac{2}{3}$$

So, $y = -\frac{2}{3}x + b$. Since the line passes

through $(2, -1)$, substitute 2 for x and -1 for y to solve for b .

$$-1 = -\frac{2}{3}(2) + b$$

$$-1 = -\frac{4}{3} + b$$

$$-\frac{3}{3} = -\frac{4}{3} + b$$

$$\frac{1}{3} = b$$

So, the equation is $y = -\frac{2}{3}x + \frac{1}{3}$.

32. First, find the slope.

$$m = \frac{-3 - (-4)}{-5 - (-2)} = \frac{1}{-3} = -\frac{1}{3}$$

So, $y = -\frac{1}{3}x + b$. Since the line passes

through $(-5, -3)$, substitute -5 for x and -3 for y to solve for b .

$$-3 = -\frac{1}{3}(-5) + b$$

$$-3 = \frac{5}{3} + b$$

$$-\frac{9}{3} = \frac{5}{3} + b$$

$$-\frac{14}{3} = b$$

So, the equation is $y = -\frac{1}{3}x - \frac{14}{3}$.

34. First, find the slope.

$$m = \frac{-2 - (-2)}{-5 - 1} = \frac{0}{-6} = 0$$

A line with zero slope is a horizontal line with an equation of the form $y = b$. Substitute the y -value for $(-5, -2)$ or $(1, -2)$ to get $y = -2$.

36. First, find the slope.

$$m = \frac{-7 - (-3)}{4 - 4} = \frac{-4}{0} = \text{undefined}$$

A line with an undefined slope is a vertical line with an equation of the form $x = c$.

Substitute the x -value for $(4, -7)$ or $(4, 3)$ to get $x = 4$.

38. First, find the slope.

$$m = \frac{7.1 - (-2.3)}{-9.4 - 3.9} = \frac{7.1 + 2.3}{-13.3} = \frac{9.4}{-13.3} \approx -0.71$$

So, $y = -0.71x + b$. Since the line contains $(3.9, -2.3)$, substitute 3.9 for x and -2.3 for y to solve for b .

$$-2.3 = -0.71(3.9) + b$$

$$-2.3 = -2.769 + b$$

$$0.47 \approx b$$

So, the equation is $y = -0.71x + 0.47$.

40. First, find the slope.

$$m = \frac{-2.21 - (-7.78)}{-7.13 - (-4.99)} = \frac{5.57}{-2.14} \approx -2.60$$

So, $y = -2.60x + b$. Since the line passes through $(-7.13, -2.21)$, substitute -7.13 for x and -2.21 for y to solve for b .

$$-2.21 = -2.60(-7.13) + b$$

$$-2.21 = 18.538 + b$$

$$-20.75 \approx b$$

So, the equation is $y = -2.60x - 20.75$.

42. The slope of the given line is 4. A line parallel to the given line has a slope of 4, so

$y = 4x + b$ is the equation for the parallel line.

Substitute 1 for x and 4 for y to solve for b since the parallel line contains $(1, 4)$.

$$4 = 4(1) + b$$

$$4 = 4 + b$$

$$0 = b$$

The parallel line's equation is $y = 4x$.

44. The slope of the given line is -1 . A line parallel to the given line also has a slope of -1 , so $y = -x + b$ is the equation for the parallel line. Substitute 2 for x and -3 for y to solve for b since the parallel line contains $(2, -3)$.

$$-3 = -1(2) + b$$

$$-3 = -2 + b$$

$$-1 = b$$

The parallel line's equation is $y = -x - 1$.

46. The slope of the given line is $-\frac{2}{3}$. A line

parallel to the given line also has a slope of $-\frac{2}{3}$, so $y = -\frac{2}{3}x + b$ is the equation for the

parallel line. Substitute 6 for x and -3 for y to solve for b since the parallel line contains $(6, -3)$.

$$-3 = -\frac{2}{3}(6) + b$$

$$-3 = -4 + b$$

$$1 = b$$

The parallel line's equation is $y = -\frac{2}{3}x + 1$.

48. To find the slope, isolate y first.

$$\begin{aligned} 5x + 2y &= 10 \\ 2y &= -5x + 10 \\ y &= -\frac{5}{2}x + 10 \end{aligned}$$

The slope is $-\frac{5}{2}$. A line parallel to the given line has the same slope. The equation for the parallel line is $y = -\frac{5}{2}x + b$. Substitute 4 for x and -1 for y to solve for b since the parallel line contains $(4, -1)$.

$$\begin{aligned} -1 &= -\frac{5}{2}(4) + b \\ -1 &= -10 + b \\ 9 &= b \end{aligned}$$

The parallel line's equation is $y = -\frac{5}{2}x + 9$.

50. To find the slope, isolate y first.

$$\begin{aligned} 3y + 5x &= -11 \\ 3y &= -5x - 11 \\ y &= -\frac{5}{3}x - \frac{11}{3} \end{aligned}$$

The slope is $-\frac{5}{3}$. A line parallel to the given line has the same slope. The equation for the parallel line is $y = -\frac{5}{3}x + b$. Substitute -1 for x and -4 for y to solve for b since the parallel line contains $(-1, -4)$.

$$\begin{aligned} -4 &= -\frac{5}{3}(-1) + b \\ -4 &= \frac{5}{3} + b \\ -\frac{12}{3} &= \frac{5}{3} + b \\ -\frac{17}{3} &= b \end{aligned}$$

The parallel line's equation is $y = -\frac{5}{3}x - \frac{17}{3}$.

52. The slope of the line $y = -4$ is 0. The line is horizontal. A line parallel to $y = -4$ has a slope of 0, as well. Since the parallel line contains $(3, -1)$ and the y -value is -1 at this point, then the equation of the line is $y = -1$.

54. The line $x = 1$ is vertical and has undefined slope. A line parallel to $x = 1$ is also vertical. Since vertical lines are of the form $x = a$, and

the parallel line contains $(-2, -5)$, the equation of the line is $x = -2$.

56. The slope of the given line is 5. A line perpendicular to the given line must have a slope of $-\frac{1}{5}$ and an equation of the form

$y = -\frac{1}{5}x + b$. Substitute 2 for x and 1 for y to solve for b since the line contains $(2, 1)$.

$$1 = -\frac{1}{5}(2) + b$$

$$1 = -\frac{2}{5} + b$$

$$\frac{5}{5} = -\frac{2}{5} + b$$

$$\frac{7}{5} = b$$

The equation of the line is $y = -\frac{1}{5}x + \frac{7}{5}$ or

$$y = -0.2x + 1.4.$$

58. The slope of the given line is -6 . A line perpendicular to the given line must then have a slope of $\frac{1}{6}$ and an equation of the form

$y = \frac{1}{6}x + b$. Substitute -3 for x and -2 for y to solve for b since the line contains $(-3, -2)$.

$$-2 = \frac{1}{6}(-3) + b$$

$$-2 = -\frac{1}{2} + b$$

$$-\frac{4}{2} = -\frac{1}{2} + b$$

$$-\frac{3}{2} = b$$

The equation of the line is $y = \frac{1}{6}x - \frac{3}{2}$.

60. The slope of the given line is $\frac{1}{3}$. A line perpendicular to the given line must then have a slope of -3 and an equation of the form $y = -3x + b$. Substitute 1 for x and -2 for y to solve for b since the line contains $(1, -2)$.

$$-2 = -3(1) + b$$

$$-2 = -3 + b$$

$$1 = b$$

The equation of the line is $y = -3x + 1$.

22 ISM: Intermediate Algebra

62. To find the slope of the given line, isolate y first.

$$\begin{aligned} 5x + 2y &= -9 \\ 2y &= -5x - 9 \\ y &= -\frac{5}{2}x - \frac{9}{2} \end{aligned}$$

The slope of the line is $-\frac{5}{2}$. A line that is perpendicular to this line must have a slope of $\frac{2}{5}$ with equation $y = \frac{2}{5}x + b$. Substitute 6 for x and -1 for y to solve for b since the line contains $(6, -1)$.

$$\begin{aligned} -1 &= \frac{2}{5}(6) + b \\ -1 &= \frac{12}{5} + b \\ -\frac{5}{5} &= \frac{12}{5} + b \\ -\frac{17}{5} &= b \end{aligned}$$

The equation of the line is $y = \frac{2}{5}x - \frac{17}{5}$ or $y = 0.4x - 3.4$.

64. To find the slope of the given line, isolate y first.

$$\begin{aligned} -3x - 4y &= 12 \\ -4y &= 3x + 12 \\ y &= -\frac{3}{4}x - 3 \end{aligned}$$

The slope of the line is $-\frac{3}{4}$. A line perpendicular to this line must have a slope of $\frac{4}{3}$ with equation $y = \frac{4}{3}x + b$. Substitute -1 for x and 2 for y to solve for b since the line contains $(-1, 2)$.

$$\begin{aligned} 2 &= \frac{4}{3}(-1) + b \\ 2 &= -\frac{4}{3} + b \\ \frac{6}{3} &= -\frac{4}{3} + b \\ \frac{10}{3} &= b \end{aligned}$$

The equation of the line is $y = \frac{4}{3}x + \frac{10}{3}$.

66. The slope of the equation, $x = -1$, is undefined. The graph of the equation is a vertical line. A line perpendicular to $x = -1$ is

a horizontal line with a slope of 0. Since this perpendicular line contains $(-4, -2)$ and the y -value at this point is -2 , the equation of the line is $y = -2$.

68. The slope of the equation, $y = 7$, is 0. The graph of the equation is a horizontal line. A line perpendicular to $y = 7$ is a vertical line with an undefined slope. Since this perpendicular line contains $(1, -1)$ and the x -value at this point is 1, the equation of the line is $x = 1$.

70. Choose any two points to find the slope.

$$m = \frac{58 - 55}{1 - 0} = \frac{3}{1} = 3$$

So, $y = 3x + b$. Since the point $(0, 55)$ is a solution to the equation, substitute 0 for x and 55 for y to solve for b .

$$\begin{aligned} 55 &= 3(0) + b \\ 55 &= 0 + b \\ 55 &= b \end{aligned}$$

The equation describing the relationship between x and y is $y = 3x + 55$.

72. Choose two points on the line to find the slope, such as $(1, 1)$ and $(4, -1)$.

$$m = \frac{1 - (-1)}{1 - 4} = \frac{1 + 1}{-3} = -\frac{2}{3}$$

So, $y = -\frac{2}{3}x + b$. Since the line contains $(1, 1)$, substitute 1 for x and 1 for y to solve for b .

$$\begin{aligned} 1 &= -\frac{2}{3}(1) + b \\ \frac{3}{3} &= -\frac{2}{3} + b \\ \frac{5}{3} &= b \end{aligned}$$

The equation for the line is $y = -\frac{2}{3}x + \frac{5}{3}$.

74. Choose two points on the line to find the slope, such as $(-2, 4)$ and $(-4, -2)$.

$$m = \frac{4 - (-2)}{-2 - (-4)} = \frac{6}{2} = 3$$

So, $y = 3x + b$. Since the line contains $(-2, 4)$, substitute -2 for x and 4 for y to solve for b .

$$4 = 3(-2) + b$$

$$4 = -6 + b$$

$$10 = b$$

The equation for the line is $y = 3x + 10$.

- 76. a.** Answers may vary. Example:
It is possible for a line to have no y -intercepts. Vertical lines of the form $x = k$ (where k is a constant not equal to 0) have no y -intercepts.
- b.** Answers may vary. Example:
It is possible for a line to have exactly one y -intercept. One example is $y = x + 1$ where the y -intercept is $(0, 1)$. In general, any equation of the form $y = mx + b$ (where m is any real number) will have exactly one y -intercept.
- c.** Answers may vary. Example:
It is not possible for a line to have exactly two y -intercepts. A line can never intersect the y -axis at exactly two points.
- d.** Answers may vary. Example:
It is possible for a line to have an infinite number of y -intercepts. The line $x = 0$ is a vertical line that lies on the y -axis and therefore, intersects it at an infinite number of points.

- 78.** No, there is no line that contains all the points. Answers may vary. Example:
The slope of the line that contains $(-1, 5)$, $(1, 1)$, $(3, -3)$ and $(4, -5)$ is -2 . So, $y = -2x + b$. Substitute -1 for x and 5 for y since the line contains $(-1, 5)$ and solve for b .
 $5 = -2(-1) + b$
 $5 = 2 + b$
 $3 = b$
The equation of the line that contains all the points (except $(-3, 7)$) is $y = -2x + 3$.

- 80.** The y values must increase by 0.5 . Therefore, the resulting line that is actually parallel to the given line has an equation of
 $y = -4x + (3 + 0.5) = -4x + 3.5$.

- 82.** Answers may vary. Example:
Find a line that satisfies $(x, y) = (3, 1)$.
For example $y = \frac{1}{3}x$. A line perpendicular to it will have slope -3 , and it should also satisfy

$$(x, y) = (3, 1)$$

$$y = -3x + b$$

$$1 = -3(3) + b$$

$$b = 10$$

$$\text{So } y = -3x + 10$$

- 84.** Answers may vary. Example:
A line with undefined slope is a vertical line. The x -coordinates of the two given points are the same.
- 86.** The equation does not include both points, although it does contain the point $(1, 5)$. Answers may vary. Example:
First find the slope.
 $m = \frac{5 - 9}{1 - 3} = \frac{-4}{-2} = 2$
So, $y = 2x + b$. Since the line contains $(1, 5)$, substitute 1 for x and 5 for y to solve for b .
 $y = 2x + b$
 $5 = 2(1) + b$
 $3 = b$
The correct equation is $y = 2x + 3$.

Homework 1.6

- 2. a.** Relation 3 and 4 could possibly be functions since each input yields only one output.
- b.** Relation 3 could be a linear function since the values of y change at a constant rate of -5 when x increases by 1 .
- 4.** Yes, this relation could be a function since each input yields exactly one output.
- 6.** No, this relation is not a function since the input, $x = 4$, yields *two* outputs, $y = 5$ and $y = 9$.
- 8.** This graph is not a function since a vertical line can intersect the graph at more than one point.
- 10.** This graph is a function since it passes the vertical line test.
- 12.** This graph is a function since it passes the vertical line test.
- 14.** This graph is not a function since a vertical line intersects it at an infinite number of points.

24 ISM: Intermediate Algebra

16. The relation $y = -3x + 8$ is a function since it can be put into the form $y = mx + b$, which defines a linear function.

18. First, isolate y .

$$\begin{aligned} 4x + 3y &= 24 \\ 3y &= -4x + 24 \\ y &= -\frac{4}{3}x + 8 \end{aligned}$$

This relation is a function since it can be put into the form $y = mx + b$ which defines a linear function.

20. $y = -1$ is a horizontal line. Because every horizontal line passes the vertical line test, $y = -1$ is a function.

22. $x = 0$ is a vertical line and does not pass the vertical line test. This is not a function.

24. First, isolate y .

$$\begin{aligned} 2x + 5y &= 9 - 4(x + 2y) \\ 2x + 5y &= 9 - 4x - 8y \\ 5y + 8y &= -2x - 4x + 9 \\ \frac{13y}{13} &= \frac{-6x + 9}{13} \\ y &= \frac{-6}{13}x + \frac{9}{13} \end{aligned}$$


This relation is a function since it can be put into the form $y = mx + b$ which defines a linear function.

26. No, a vertical line is not a function since it does not pass the vertical line test.

28. Yes, a semicircle that is the “upper half” of a circle is a function since no vertical line intersects the semicircle at more than one point.

30. a.

x	y
-4	$\frac{1}{2}(-4) + 2 = 0$
-2	$\frac{1}{2}(-2) + 2 = 1$
0	$\frac{1}{2}(0) + 2 = 2$
2	$\frac{1}{2}(2) + 2 = 3$
4	$\frac{1}{2}(4) + 2 = 4$



c. For each input–output pair, the output is 2 more than $\frac{1}{2}$ times the input.

32. The domain is $-2 \leq x \leq 3$ and the range is $-2 \leq y \leq 4$.

34. The domain is $-4 \leq x \leq 5$ and the range is $-3 \leq y \leq 4$. The highest point on the graph appears to include a y -value of 4.


36. The domain is $-3 \leq x \leq 5$ and the range is $0 \leq y \leq 4$. The lowest point on the graph appears to include a y -value of 0 and the highest point appears to include 4.

38. The domain is $-3 \leq x \leq 3$ and the range is $0 \leq y \leq 3$.

40. The domain is all real numbers and the range is $y \geq 1$.


42. The domain is all real numbers and the range is all real numbers.

44. Sketch the graph of $y = x^4$.




Note that since this graph passes the vertical line test, $y = x^4$ is a function.

46. Sketch the graph of $y^3 = x$.



Note that since this graph passes the vertical line test, $y^3 = x$ is a function.

48. Answers may vary. Example:




x	y
-3	2
0	2
3	2
3	-2
-3	-2

The rectangle relation is not a function since a vertical line can intersect the graph of the rectangle at more than one point.


50. Answers may vary. Example:

Suppose that when $x = -4$, $y = 0, 1$, and 2 . Then suppose when $x = 5$, $y = 2$. Sketch the graph.




This relation is not a function since it does not pass the vertical line test.

52. Answers may vary. Example:



54. Answers may vary. Example:




56. Answers may vary. Example:

If a vertical line intersects the graph of a relation more than once, then at least one input to the relation must have two or more outputs, and therefore the relation is not a function.


58. Answers may vary.

Chapter 1 Review Exercises

- The amount of time it takes to read and reply to texts depends on the number of texts that a person receives. The amount of time to read and reply (t) is the response variable and the number of texts (n) is the explanatory variable.
- The length of the candle (L) is the response variable and minutes (t) is the explanatory variable.



- The amount of time (T) to cook a marshmallow is the response variable and the distance (d) from the campfire is the explanatory variable.



$$\begin{aligned}
 4. \quad & 3(2x-4)-2=5x-(3-4x) \\
 & 6x-12-2=5x-3+4x \\
 & 6x-14=9x-3 \\
 & \frac{-3x}{-3}=\frac{11}{-3} \\
 & x=-\frac{11}{3}
 \end{aligned}$$

$$\begin{aligned}
 5. \quad & \frac{2}{3}w-\frac{1}{2}=\frac{5}{6}w+\frac{4}{3} \\
 6\left(\frac{2}{3}w-\frac{1}{2}\right) &= 6\left(\frac{5}{6}w+\frac{4}{3}\right) \\
 4w-3 &= 5w+8 \\
 -11 &= w
 \end{aligned}$$

$$\begin{aligned}
 6. \quad & a(x-c)=d \\
 x-c &= \frac{d}{a} \\
 x &= \frac{d}{a}+c
 \end{aligned}$$

$$\begin{aligned}
 7. \quad & \text{To find the } x\text{-intercept, substitute } 0 \text{ for } y. \\
 & 3x-5y=17 \\
 & 3x-5(0)=17 \\
 & \frac{3x}{3}=\frac{17}{3} \\
 & x=\frac{17}{3}
 \end{aligned}$$

The x -intercept is $\left(\frac{17}{3}, 0\right)$.

To find the y -intercept, substitute 0 for x .

$$\begin{aligned}
 & 3x-5y=17 \\
 & 3(0)-5y=17 \\
 & \frac{-5y}{-5}=\frac{17}{-5} \\
 & y=-\frac{17}{5}
 \end{aligned}$$

The y -intercept is $\left(0, -\frac{17}{5}\right)$

8. To find the x -intercept, substitute 0 for y .

$$\begin{aligned}
 ax+b &= cy \\
 ax+b &= (0)y \\
 \frac{ax}{a} &= \frac{-b}{a} \\
 x &= -\frac{b}{a}
 \end{aligned}$$

The x -intercept is $\left(-\frac{b}{a}, 0\right)$.

To find the y -intercept, substitute 0 for x .

$$\begin{aligned}
 ax+b &= cy \\
 (0)x+b &= cy \\
 \frac{b}{c} &= y
 \end{aligned}$$

The y -intercept is $\left(0, \frac{b}{c}\right)$

$$9. \quad y = -4$$

$$10. \quad y \approx -\frac{2}{3}$$

$$11. \quad x = -4$$

$$12. \quad x = -1$$

$$13. \quad m = \frac{-2-(-5)}{-3-2} = \frac{3}{-5} = -\frac{3}{5}$$

The line is decreasing.

$$14. \quad m = \frac{-7-(-3)}{-9-(-1)} = \frac{-4}{-8} = \frac{1}{2}$$

The line is increasing.


$$15. \quad m = \frac{-1-3}{4-4} = \frac{-4}{0} \text{ is undefined}$$

The line is vertical.

$$16. \quad m = \frac{2.99-(-8.48)}{-5.27-3.54} = \frac{11.47}{-8.81} \approx -1.30$$

The line with slope of -1.30 is decreasing.

17. Use the slope, -3 , and the y -intercept, 10 , to graph the equation.




18. First, isolate y .


$$y + 2x = 0$$

$$y = -2x$$

Use the slope, -2 , and the y -intercept, 0 , to graph the equation.



19. $y = 7$ is a horizontal line with a y -intercept of 7 .




20. First, isolate y .

$$3x - 2y = 12$$

$$-2y = -3x + 12$$

$$y = \frac{3}{2}x - 6$$

Use the slope $\frac{3}{2}$, and the y -intercept, -6 , to graph the equation.



21. First, isolate y .


$$-3(y + 2) = 2x + 9$$

$$-3y - 6 = 2x + 9$$

$$-3y = 2x + 15$$

$$y = -\frac{2}{3}x - 5$$

Use the slope, $-\frac{2}{3}$, and the y -intercept, -5 , to graph the equation.



22. First, isolate y .


$$3x - 2(2y - 1) = 8x - 3(x + 2)$$

$$3x - 4y + 2 = 8x - 3x - 6$$

$$\frac{-4y}{-4} = \frac{2x - 8}{-4}$$

$$y = -\frac{1}{2}x + 2$$

Use the slope, $-\frac{1}{2}$, and the y -intercept, 2 , to graph the equation.



23. First, isolate y .

$$a(x - y) = c$$

$$x - y = \frac{c}{a}$$

$$-y = -x + \frac{c}{a}$$

$$y = x - \frac{c}{a}$$

The slope is 1 and the y -intercept is $(0, -\frac{c}{a})$.

24. Solve the first equation for y to find the slope.

$$2x + 5y = 7$$

$$5y = -2x + 7$$

$$y = -\frac{2}{5}x + \frac{7}{5}$$

The slopes are $-\frac{2}{5}$ and $\frac{2}{5}$. They are not

reciprocals, although they are additive inverses of each other. The lines are neither parallel nor perpendicular.

25. Solve each equation for y to find the slope.

$$3x - 8y = 7 \qquad -6x + 16y = 5$$

$$\frac{-8y}{-8} = \frac{-3x + 7}{-8} \qquad \frac{16y}{16} = \frac{6x + 5}{16}$$

$$y = \frac{3}{8}x - \frac{7}{8} \qquad y = \frac{3}{8}x + \frac{5}{16}$$

The slopes are $\frac{3}{8}$ and $\frac{3}{8}$. The lines are parallel.

26. We are given the slope, $m = -4$, and a point on the line, $(-3, 7)$. Use $y = mx + b$ to find b .

$$\begin{aligned}y &= mx + b \\7 &= -4(-3) + b \\7 &= 12 + b \\-5 &= b\end{aligned}$$

Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$\begin{aligned}y &= -4x + (-5) \\y &= -4x - 5\end{aligned}$$

27. We are given the slope, $m = -\frac{2}{3}$, and a point on the line, $(5, -4)$. Use $y = mx + b$ to find b .

$$\begin{aligned}y &= mx + b \\-4 &= -\frac{2}{3}(5) + b \\-4 &= -\frac{10}{3} + b \\-\frac{12}{3} &= -\frac{10}{3} + b \\-\frac{2}{3} &= b\end{aligned}$$

Now, substitute for m and b in slope-intercept form to obtain the equation of the line.

$$\begin{aligned}y &= -\frac{2}{3}x + \left(-\frac{2}{3}\right) \\y &= -\frac{2}{3}x - \frac{2}{3}\end{aligned}$$

28. First, find the slope.

$$m = \frac{6 - (-2)}{2 - (-3)} = \frac{8}{5}$$

So, $y = \frac{8}{5}x + b$. Since the line contains $(2, 6)$, substitute 2 for x and 6 for y and solve for b .

$$\begin{aligned}6 &= \frac{8}{5}(2) + b \\6 &= \frac{16}{5} + b \\ \frac{30}{5} &= \frac{16}{5} + b \\ \frac{14}{5} &= b\end{aligned}$$

So, the equation is $y = \frac{8}{5}x + \frac{14}{5}$.

29. First, find the slope.

$$m = \frac{6 - (-2)}{-4 - 2} = \frac{8}{-6} = -\frac{4}{3}$$

So, $y = -\frac{4}{3}x + b$. Since the line contains

$(2, -2)$, substitute 2 for x and -2 for y and solve for b .

$$\begin{aligned}-2 &= -\frac{4}{3}(2) + b \\-\frac{6}{3} &= -\frac{8}{3} + b \\ \frac{2}{3} &= b\end{aligned}$$

So, the equation is $y = -\frac{4}{3}x + \frac{2}{3}$.

30. First, find the slope.

$$m = \frac{5 - (-2)}{3 - 3} = \frac{7}{0} \text{ undefined}$$

A line with undefined slope is a vertical line of the form $x = a$. Since $(3, 5)$ is a point on the line and the value of x is 3 at that point, the equation of the line is $x = 3$.

31. First, find the slope.

$$m = \frac{-8.79 - (-6.38)}{-3.62 - 2.51} = \frac{-2.41}{-6.13} \approx 0.39$$

So, $y = 0.39x + b$. Since the line contains $(-3.62, -8.79)$, substitute -3.62 for x and -8.79 for y and solve for b .

$$\begin{aligned}y &= 0.39x + b \\-8.79 &= 0.39(-3.62) + b \\-8.79 &= -1.4118 + b \\-7.37 &\approx b\end{aligned}$$

So, the equation is $y = 0.39x - 7.37$.

32. Line 1:

Two points on line 1 are $(1, -2)$ and $(-1, 2)$. Use these points to find the slope.

$$m = \frac{2 - (-2)}{-1 - 1} = \frac{4}{-2} = -2$$

So, $y = -2x + b$. We can see from the graph that the y -intercept is 0. So, the equation of the line is $y = -2x + 0 = -2x$.

Line 2:

Two points on line 1 are $(-1, 0)$ and $(0, 2)$. Use these points to find the slope.

$$m = \frac{2 - 0}{0 - (-1)} = \frac{2}{1} = 2$$

So, $y = 2x + b$. We can see from the graph that the y -intercept is 2. So, the equation of the line is $y = 2x + 2$.

Line 3:

Since line 3 is a horizontal line, it is of the form $y = b$. Since the y -intercept is -3 , the equation of the line is $y = -3$.

33. The y -values decrease from 20 to 4 (for a change of -16) while the x -values increase from 2 to 6 (for a change of 4). So, the slope is $\frac{-16}{4} = \frac{-4}{1} = -4$, or the corresponding y -values decrease 4 for every x -value increase of 1. Complete the table by subtracting 4 from each y -value.

x	y
2	20
3	16
4	12
5	8
6	4
7	0

34. Line 1 is parallel to $y = 0.5x + 6$. Since parallel lines have the same slope, the slope of line 1 is 0.5. The equation of line 1 is $y = 0.5x + b$. Line 1 has the same y -intercept as the line $y = -1.5x + 3$. The y -intercept is $b = 3$. The equation of line 1 is $y = 0.5x + 3$.

35. a. B, C and F
 b. C and E
 c. C
 d. A and D

36. Since the line is parallel to the line $3x - y = 6$, or $y = 3x - 6$, it will have a slope of 3. Use the slope, 3, and the point on the line, $(-2, 5)$, to find the y -intercept.

$$5 = 3(-2) + b$$

$$5 = -6 + b$$

$$11 = b$$

The equation of the line is $y = 3x + 11$.

37. A line with an infinite number of x -intercepts is the line that lies on the x -axis. This is the horizontal line $y = 0$.
38. Relations 1 and 3 could possibly be functions, since there is only one output for each input. Relations 2 and 4 could not possibly be functions, since there are two or more outputs for a single input.
39. The graph is not a function because a vertical line can intersect the graph more than once.

40. First, isolate y .

$$5x - 6y = 3$$


$$\frac{-6y}{-6} = \frac{-5x + 3}{-6}$$

$$y = \frac{5}{6}x - 3$$

This relation is a function since it can be put into the form $y = mx + b$ which defines a linear function.

41. $x = 9$ is a vertical line. It does not pass the vertical line test and is not a function.

42. Sketch the graph of $y^2 = x$.




Note that this graph does not pass the vertical line test. Therefore, $y^2 = x$ is not a function.

43. The domain is all real numbers and the range is $y \leq 4$.

Chapter 1 Test

1. The value in dollars of gold depends on the weight of a gold bar. The value (v) is the response variable and the weight (w) is the explanatory variable.
2. Answers may vary. Example:



3. Answers may vary. Example: The diameter of a balloon in inches (y) is the response variable and the number of seconds after the knot of the balloon was untied (x) is the explanatory variable.