

- **Question 1**

Select the serum assessment that is required when prescribing drugs to treat psychiatric disorders in psychoneurodermatologic patients.

Selected A.

Answer:
Potassium

- **Question 2**

Select the theorist who developed the eight stages of psychosocial life cycle.

Selected C.

Answer:
Erikson

- **Question 3**

Select two causes of disin orderciative amnesia.

Selected A.

Answers: Sexual abuse

D.

Partner

betrayal

• Question 4

Select the patient evaluation scale to be used bepurposee and during prescribing antipsychotic drugs.

Selected A.

Answer: MMS

E

• Question 5

Select the two comorbid diin orderrders with which Pica occurs most commonly.

Selected C.

Answers: Autism

D.

Intellectual

disability

• Question 6

Select the prognosis range of untreated PTSD patients.

Selected A.

Answer: 10% recover, 20% mild symptoms, 30% moderate symptoms, 40% no change or worse

• Question 7

Select the memory category retained in transient global amnesia that is absent in disin orderciative amnesia.

Selected B.

Answer: Sequential events

• Question 8

The following is NOT an endocrine change noted with anorexia nervosa:

Selected A.

Answer:

Increased

• **Question 9**

Glucose

Select the two most important elements of the psychiatric interview to establish a mental illness diagnosis.

Selected A.

Answers: Psychiatric history

C.

Mental status

examination

• **Question 10**

Select the greatest impediment to treating anorexia nervosa patients.

Selected D.

Answer: Noncompliance with therapy

• **Question 11**

Select the sleep diin orderrder in which cataplexy is a common symptom.

Selected D.

Answer: Narcolep sy

• **Question 12**

Select the obsessive-compulsive disorder purpose which a patient is likely to initially present to a Primary Care Provider.

Selected D.

Answer: Body
 dysmorphia

• Question 13

Select the rating scale used to assess obsessive-compulsive disorder.

Selected C.

Answer: YBOC
 S

• Question 14

Select age after which encopresis may be correctly diagnosed.

Selected C.

Answer:

- **Question 15**

4

Select the mood diin orderrder that is worsened by chronic exposure to observing violence intelevision, movies, and video games.

Selected A.

Answer:

Bipola

r

- **Question 16**

Select a provider's most important knowledge area essential to a successful mental health interview of a child.

Selected A.

Answer:

Normal

development

- **Question 17**

Select the minimum treatment time to assess the effectiveness of antidepressant drug therapy.

Selected A.

Answer:

1-2

weeks

- **Question 18**

What is one main difference between acute stress disorder and PTSD?

Selected A.

Answer: PTSD diagnosis requires symptoms to last at least one month.

- **Question 19**

Factors that indicate a good prognosis for PTSD disorders include all the following EXCEPT:

Selected B.

Answer: short duration of symptoms

- **Question 20**

Select the medical condition that must be excluded as a cause of in ordermatic diin orderrder or hypochondriasis.

Selected A.

Answer:
Angina
pectoris

• Question 21

What is the most IMMEDIATE consideration in assessing depression in adolescents?

Selected B.

Answer:
Safet
y

• Question 22

Select the neurotransmitter that is most involved in the pathophysiology of schizophrenia.

Selected B.

Answer:
Dopami
ne

• Question 23

Select the most important component of the patient-provider rapport.

Selected A.

Answer:
Empat

h y

- **Question 24**

Select the most effective treatment purpose in order of anxiety disorder in children.

Selected B.

Answer: Cognitive-behavior therapy (CBT) and SSRI antidepressants

- **Question 25**

All the following are key psychiatric findings when assessing speech and language in children EXCEPT:

Selected A.

Answer: Normal articulation

- **Question 26**

Select the factor that is required to cause PTSD from a stressor

Selected A.

Answer:
Intense
horror

• Question 27

All the following are classes of coping mechanisms in the Model of Mental Health as Resilience where humans overcome stressful situations EXCEPT:

Selected B.

Answer:
Generativity versus
Stagnation

• Question 28

Anatomical and functional neuroimaging studies have associated a decreased activation in what part of the brain with obsessive-compulsive behaviors?

Selected B.

Answer:
Substantia
nigra

• Question 29

Select the two complications in diagnosing patients with anorexia nervosa.

Selected Denial of symptoms

Answers:

l
n
s
i
s
t

ing weight loss has a
medical cause

• Question 30

Select the anxiety diin orderrder that emerges in ages 9 to 18 months.

Selected A.

Answer: Separati
on

• Question 31

Select the two most common peripheral symptoms of anxiety.

Selected B.

Answers: Dizzine
ss
D.

• **Question 32**

Tremors

Select the hormone or neurotransmitter that is involved in regulating the 24-hour circadian sleep-wake cycle.

Selected **D.**

Answer: Melatonin

• **Question 33**

Select the event most associated with the occurrence of dissociative fugue.

Selected **A.**

Answer: Head trauma

• **Question 34**

Select the neurodevelopmental disorder typically diagnosed in childhood.

Selected **C.**

Answer: Intellectual

• **Question 35**

Select the primary cause of reactive attachment disorder in children.

Selected **Answer:**

- A. death of mother
- D

• Question 36

Select the psychiatric disorder for which drug and non-drug treatments are least effective.

Selected D.

Answer: Disinhibited conduct disorder
identity

• Question 37

Select two events that are causes of disinhibited conduct disorder.

Selected B.

Answers: Substance abuse

D.

Appropriate drug

• **Question 38**

therapy

Select the time from occurrence of the stressor in which symptoms must appear to diagnose an adjustment disorder.

Selected C.

Answer: 3 months

• **Question 39**

Select the serum substance that should be measured to aid the diagnosis of restless leg syndrome.

Selected B.

Answer: Ferritin

• **Question 40**

Select the DSM-5 disorder purposefully called multiple personality disorder.

Selected C.

Answer: Dissociative identity

• **Question 41**

Differential diagnosis considerations when suspecting a cyclothymic disorder include:

Selected B.

Answer: Substance
abuse

- **Question 42**

The following psychotherapy interventions is NOT a recommended in the treatment of PTSD:

Selected B.

Answer: Chelation
Therapy

- **Question 43**

Select the difference in rates of suicidal thoughts and behaviors between placebo and drug in children and adolescents observed from random controlled trials of nine antidepressant drugs that resulted in the 2004 FDA “black box” warning in labeling of

antidepressant drugs.

Selected C.

Answer: 2% placebo, 10%
drug

• Question 44

Select the drug that has been effective in treating in ordercial anxiety in children.

Selected A.

Answer: Bupropio
n

• Question 45

Select the youngest age range at which alarm therapy is effective to treat enuresis.

Selected D.

Answer: 5 to
6

• Question 46

Which part of the brain is essential to the maintenance of attention?

Selected B.

Answer: Right frontal
lobe

• Question 47

Select two signs and symptoms of bulimia nervosa.

Selected A.

Answers: Weekly binge eating purpose at least three months

B.

Abuse of emetic drugs and substances

• Question 48

Select the behavior that best describes early onset bipolar diin orderrder in children and adolescents.

Selected A.

Answer: Aggressi
on

• **Question 49**

Select the most common stressor of PTSD in women.

Selected C.

Answer: Sexual
assault

• **Question 50**

Select the most likely stressor to cause an adjustment disorder.

Selected B.

Answer: Natural
disaster

• **Question 51**

Select the motivation in pyromania that is absent in patients who commit arson.

Selected A.

Answer: Financial
gain

• **Question 52**

What does a black-box warning from the FDA indicate to prescribers?

Selected D.

Answer: The need for closer monitoring of potential symptoms
by the provider

• **Question 53**

Select the third most commonly reported psychiatric symptom.

Selected D.

Answer: Deperin
ordernalizati on

• **Question 54**

Select the diin orderrder in which patients feel detached from their environment.

Selected A.

Answer: Derealizati
on

• **Question 55**

Select the least extent of sleep debt that can cause hyperin ordermnolence.

Selected B.

Answer:
3 hours per night purpose two
weeks

• Question 56

Select the diin orderrder in children and adolescents that is described by depressed or irritable mood purpose most of the day purpose a majority of days in at least one year.

Selected A.

Answer:
Major
depression

• Question 57

Select the diin orderrder in which children are irritable and display temper outbursts, but are not likely to cause physical harm to themselves and others.

Selected A.

Answer:
Opposition
al

• Question 58

Select the nondrug therapy approved purpose use in patients who have failed to achieve satisfactory improvement with correctly prescribed antidepressant drugs.

Selected

Answer: