
CONTENTS

Chapter 1.1	Propositions and Connectives	1
1.2	Conditionals and Biconditionals	6
1.3	Quantifiers	13
1.4	Basic Proof Methods I	17
1.5	Basic Proof Methods II	22
1.6	Proofs Involving Quantifiers	26
1.7	Additional Examples of Proofs	29
Chapter 2.1	Basic Concepts of Set Theory	38
2.2	Set Operations	40
2.3	Extended Set Operations and Indexed Families of Sets	46
2.4	Mathematical Induction	49
2.5	Equivalent Forms of Induction	59
2.6	Principles of Counting	62
Chapter 3.1	Cartesian Products and Relations	67
3.2	Equivalence Relations	70
3.3	Partitions	75
3.4	Ordering Relations	78
3.5	Graphs	82
Chapter 4.1	Functions as Relations	85
4.2	Constructions of Functions	88
4.3	Functions That Are Onto; One-to-One Functions	92
4.4	One-to-One Correspondences and Inverse Functions	95
4.5	Images of Sets	199
4.6	Sequences	102
Chapter 5.1	Equivalent Sets; Finite Sets	105
5.2	Infinite Sets	108
5.3	Countable Sets	111
5.4	The Ordering of Cardinal Numbers	114
5.5	Comparability of Cardinal Numbers and the Axiom of Choice	117
Chapter 6.1	Algebraic Structures	119
6.2	Groups	122
6.3	Subgroups	126
6.4	Operation Preserving Maps	128
6.5	Rings and Fields	131
Chapter 7.1	Completeness of the Real Numbers	133
7.2	The Heine-Borel Theorem	136
7.3	The Bolzano-Weierstrass Theorem	139
7.4	The Bounded Monotone Sequence Theorem	140
7.5	Equivalents of Completeness	143

1 Logic and Proofs

1.1 Propositions and Connectives

1. (a) true (b) false (c) true (d) false
 (e) false (f) false (g) false (h) false
2. (a) Not a proposition
 (b) False proposition
 (c) Not a proposition. It would be a proposition if a value for x had been assigned.
 (d) Not a proposition. It would be a proposition if values for x and y had been assigned.
 (e) False proposition
 (f) True proposition
 (g) False proposition
 (h) True proposition
 (i) False proposition
 (j) Not a proposition. It is neither true nor false.

3. (a)

P	$\sim P$	$P \wedge \sim P$
T	F	T
F	T	F

(b)

P	$\sim P$	$P \vee \sim P$
T	F	T
F	T	T

(c)

P	Q	$\sim Q$	$P \wedge \sim Q$
T	T	F	F
F	T	F	F
T	F	T	T
F	F	T	F

(d)

P	Q	$\sim Q$	$Q \vee \sim Q$	$P \wedge (Q \vee \sim Q)$
T	T	F	T	T
F	T	F	T	F
T	F	T	T	T
F	F	T	T	F

(e)

P	Q	$\sim Q$	$P \wedge Q$	$(P \wedge Q) \vee \sim Q$
T	T	F	T	T
F	T	F	F	F
T	F	T	F	T
F	F	T	F	T

(f)

P	Q	$P \wedge Q$	$\sim (P \wedge Q)$
T	T	T	F
F	T	F	T
T	F	F	T
F	F	F	T

(g)

P	Q	R	$\sim Q$	$P \vee \sim Q$	$(P \vee \sim Q) \wedge R$
T	T	T	F	T	T
F	T	T	F	F	F
T	F	T	T	T	T
F	F	T	T	T	T
T	T	F	F	T	F
F	T	F	F	F	F
T	F	F	T	T	F
F	F	F	T	T	F

(h)

P	Q	$\sim P$	$\sim Q$	$\sim P \wedge \sim Q$
T	T	F	F	F
F	T	T	F	F
T	F	F	T	F
F	F	T	T	T

(i)

P	Q	R	$Q \vee R$	$P \wedge (Q \vee R)$
T	T	T	T	T
F	T	T	T	F
T	F	T	T	T
F	F	T	T	F
T	T	F	T	T
F	T	F	T	F
T	F	F	F	F
F	F	F	F	F

(j)

P	Q	R	$P \wedge Q$	$P \wedge R$	$(P \wedge Q) \vee (P \wedge R)$
T	T	T	T	T	T
F	T	T	F	F	F
T	F	T	F	T	T
F	F	T	F	F	F
T	T	F	T	F	T
F	T	F	F	F	F
T	F	F	F	F	F
F	F	F	F	F	F

4. (a) false (b) true (c) true (d) true
 (e) false (f) false (g) false (h) false
 (i) true (j) true (k) false (l) false

5. (a) No solution.

(b)

P	Q	$P \vee Q$	$Q \vee P$
T	T	T	T
F	T	T	T
T	F	T	T
F	F	F	F

Since the third and fourth columns are the same, the propositions are equivalent.

Since the fifth and eighth columns are the same, the propositions are equivalent.

(h) No solution.

(i)

P	Q	$P \vee Q$	$\sim (P \vee Q)$	$\sim P$	$\sim Q$	$\sim P \wedge \sim Q$
T	T	T	F	F	F	F
F	T	T	F	T	F	F
T	F	T	F	F	T	F
F	F	F	T	T	T	T

Since the fourth and eighth columns are the same, the propositions are equivalent.

6. (a) equivalent (b) equivalent
 (c) equivalent (d) equivalent
 (e) equivalent (f) not equivalent
 (g) not equivalent (h) not equivalent
7. (a) $\sim P$, true (b) $P \wedge Q$, true
 (c) $P \vee Q$, true (d) $P \vee Q \vee R$, true
8. (a) Since P is equivalent to Q , P has the same truth table as Q . Therefore, Q has the same truth table as P , so Q is equivalent to P .
 (b) Since P is equivalent to Q , P and Q have the same truth table. Since Q is equivalent to R , Q and R have the same truth table. Thus, P and R have the same truth table so P is equivalent to R .
 (c) Since P is equivalent to Q , P and Q have the same truth table. That is, the truth table for P has value true on exactly the same lines that the truth table for Q has value true. Therefore the truth table for $\sim Q$ has value false on exactly the same lines that the truth table for $\sim P$ has the value false. Thus $\sim Q$ and $\sim P$ have the same truth table.
9. (a) $(P \wedge Q) \vee (\sim P \wedge \sim Q)$ is neither.

P	Q	$\sim P$	$\sim Q$	$P \wedge Q$	$\sim P \wedge \sim Q$	$(P \wedge Q) \vee (\sim P \wedge \sim Q)$
T	T	F	F	T	F	T
F	T	T	F	F	F	F
T	F	F	T	F	F	F
F	F	T	T	F	T	T

(b) $\sim (P \wedge \sim P)$ is a tautology.

P	$\sim P$	$P \wedge \sim P$	$\sim (P \wedge \sim P)$
T	F	F	T
F	T	F	T

(c) $(P \wedge Q) \vee (\sim P \vee \sim Q)$ is a tautology.

P	Q	$\sim P$	$\sim Q$	$P \wedge Q$	$\sim P \vee \sim Q$	$(P \wedge Q) \vee (\sim P \vee \sim Q)$
T	T	F	F	T	F	T
F	T	T	F	F	T	T
T	F	F	T	F	T	T
F	F	T	T	F	T	T

(d) $(A \wedge B) \vee (A \wedge \sim B) \vee (\sim A \wedge B) \vee (\sim A \wedge \sim B)$ is a tautology.

A	B	$\sim A$	$\sim B$	$A \wedge B$	$A \wedge \sim B$	$\sim A \wedge B$	$\sim A \wedge \sim B$	$(A \wedge B) \vee (A \wedge \sim B) \vee (\sim A \wedge B) \vee (\sim A \wedge \sim B)$
T	T	F	F	T	F	F	F	T
F	T	T	F	F	F	T	F	T
T	F	F	T	F	T	F	F	T
F	F	T	T	F	F	F	T	T

(e) $(Q \wedge \sim P) \wedge \sim (P \wedge R)$ is neither.

P	Q	R	$\sim P$	$Q \wedge \sim P$	$P \wedge R$	$\sim (P \wedge R)$	$(Q \wedge \sim P) \wedge \sim (P \wedge R)$
T	T	T	F	F	T	F	F
F	T	T	T	T	F	T	T
T	F	T	F	F	T	F	F
F	F	T	T	F	F	T	F
T	T	F	F	F	F	T	F
F	T	F	T	T	F	T	T
T	F	F	F	F	F	T	F
F	F	F	T	F	F	T	F

(f) $P \vee [(\sim Q \wedge P) \wedge (R \vee Q)]$ is neither.

P	Q	R	$\sim Q$	$\sim Q \wedge P$	$R \vee Q$	$[(\sim Q \wedge P) \wedge (R \vee Q)]$	$P \vee [(\sim Q \wedge P) \wedge (R \vee Q)]$
T	T	T	F	F	T	F	T
F	T	T	F	F	T	F	F
T	F	T	T	T	T	T	T
F	F	T	T	F	T	F	F
T	T	F	F	F	T	F	T
F	T	F	F	F	T	F	F
T	F	F	T	T	F	F	T
F	F	F	T	F	F	F	F

10. (a) contradiction (b) tautology
(c) tautology (d) tautology
11. (a) x is not a positive integer.
(b) Cleveland will lose the first game and the second game. Or, Cleveland will lose both games.
(c) $5 < 3$
(d) 641,371 is not composite. Or 641,371 is prime.
(e) Roses are not red or violets are not blue.
(f) T is bounded and T is not compact.
(g) M is not odd or M is not one-to-one.
(h) The function f does not have a positive first derivative at x or does not have a positive second derivative at x .
(i) The function g does not have a relative maximum at $x = 2$ (deleted comma) and does not have a relative maximum at $x = 4$, or else g does not have a relative minimum at $x = 3$.
(j) $z < s$ or $z \leq t$.
(k) R is not transitive or R is reflexive.
(l) If the function g has a relative minimum at $x = 2$ or $x = 4$, then g does not have a relative minimum at $x = 3$.
12. (a) $[\sim(\sim P)] \vee [(\sim Q) \wedge (\sim S)]$
(b) $[Q \wedge (\sim S)] \vee \sim(P \wedge [Q \wedge (\sim S)]) \vee \sim((\sim P \wedge Q))$.
(c) $[[P \wedge (\sim Q)] \vee [(\sim P) \wedge (\sim R)]] \vee [(\sim P) \wedge S]$
(d) $[(\sim P) \vee ([Q \wedge (\sim P)]) \wedge Q] \vee R$.

13. (a) i.

A	B	$A \oplus B$
T	T	F
F	T	T
T	F	T
F	F	F

ii.

A	B	$A \vee B$	$A \wedge B$	$\sim (A \wedge B)$	$(A \vee B) \wedge \sim (A \wedge B)$
T	T	T	T	F	F
F	T	T	F	T	T
T	F	T	F	T	T
F	F	F	F	T	F

Since the final columns of the two tables are identical, the two propositions have the same truth table, thus they are equivalent.

(b) i.

A	B	$A \text{ NAND } B$	$A \text{ NOR } B$
T	T	F	F
F	T	T	F
T	F	T	F
F	F	T	T

ii.

A	B	$A \text{ NAND } B$	$A \text{ NOR } B$	$(A \text{ NAND } B) \vee (A \text{ NOR } B)$
T	T	F	F	F
F	T	T	F	T
T	F	T	F	T
F	F	T	T	T

Since the third and last columns are equal, the propositions are equivalent.

iii.

A	B	$A \text{ NAND } B$	$A \text{ NOR } B$	$(A \text{ NAND } B) \wedge (A \text{ NOR } B)$
T	T	F	F	F
F	T	T	F	F
T	F	T	F	F
F	F	T	T	T

Since the fourth and last columns are equal, the propositions are equivalent.

1.2 Conditionals and Biconditionals

1. (a) Antecedent: squares have three.
Consequent: triangles have four sides.
- (b) Antecedent: The moon is made of cheese.
Consequent: 8 is an irrational number.
- (c) Antecedent: b divides 3.
Consequent: b divides 9.
- (d) Antecedent: f is differentiable.
Consequent: f is continuous.
- (e) Antecedent: a is convergent.
Consequent: a is bounded.
- (f) Antecedent: f if integrable.
Consequent: f is bounded.
- (g) Antecedent: $1 + 1 = 2$.
Consequent: $1 + 2 = 3$.

- (h) Antecedent: the fish bite.
Consequent: the moon is full.
- (i) Antecedent: An athlete qualifies for the Olympic team.
Consequent: The athlete has a time of 3 minutes, 48 seconds or less (in the event).
2. (a) Converse: If triangles have four sides, then squares have three sides.
Contrapositive: If triangles do not have four sides, then squares do not have three sides.
- (b) Converse: If 8 is irrational, then the moon is made of cheese.
Contrapositive: If 8 is rational, then the moon is not made of cheese.
- (c) Converse: If b divides 9, then b divides 3.
Contrapositive: If b does not divide 9, then b does not divide 3.
- (d) Converse: If f is continuous, then f is differentiable.
Contrapositive: If f is not continuous, then f is not differentiable.
- (e) Converse: If a is bounded, then a is convergent.
Contrapositive: If a is not bounded, then a is not convergent.
- (f) Converse: If f is bounded, then f is integrable.
Contrapositive: If f is not bounded, then f is not integrable.
- (g) Converse: If $1 + 2 = 3$, then $1 + 1 = 2$.
Contrapositive: If $1 + 1 \neq 2$, then $1 + 2 \neq 3$.
- (h) Converse: If the moon is full, then fish will bite.
Contrapositive: If the moon is not full, then fish will not bite.
- (i) Converse: A time of 3 minutes, 48 seconds or less is sufficient to qualify for the Olympic team.
Contrapositive: If an athlete records a time that is not 3 minutes and 48 seconds or less, then that athlete does not qualify for the Olympic team.
3. (a) Q may be either true or false.
(b) Q must be true.
(c) Q must be false.
(d) Q must be false.
(e) Q must be false.
4. (a) Antecedent: $A(x)$ is an open sentence with variable x .
Consequent: $\sim (\forall x)A(x)$ is equivalent to $(\exists x) \sim A(x)$.
- (b) Antecedent: Every even natural number greater than 2 is the sum of two primes.
Consequent: Every odd natural number greater than 5 is the sum of three primes.
- (c) Antecedent: A is a set with n elements.
Consequent: $\mathcal{P}(A)$ is a set with 2^n elements.
- (d) Antecedent: S is a subset of \mathbb{N} such that $1 \in S$ and, for all $n \in \mathbb{N}$, if $n \in S$, then $n + 1 \in S$.
Consequent: $S = \mathbb{N}$.
- (e) Antecedent: A is a finite set with m elements and B is a finite set with n elements.
Consequent: $\overline{A \times B} = mn$.

(f)

P	Q	R	S	$Q \Rightarrow S$	$Q \Rightarrow R$	$P \vee Q$	$S \vee R$	$(Q \Rightarrow S) \wedge (Q \Rightarrow R)$
T	T	T	T	T	T	T	T	T
F	T	T	T	T	T	T	T	T
T	F	T	T	T	T	T	T	T
F	F	T	T	T	T	F	T	T
T	T	F	T	T	F	T	T	F
F	T	F	T	T	F	T	T	F
T	F	F	T	T	T	T	T	T
F	F	F	T	T	T	F	T	T
T	T	T	F	F	T	T	T	F
F	T	T	F	F	T	T	T	F
T	F	T	F	T	T	T	T	T
F	F	T	F	T	T	F	T	T
T	T	F	F	F	F	T	F	F
F	T	F	F	F	F	T	F	F
T	F	F	F	T	T	T	F	T
F	F	F	F	T	T	F	F	T

P	Q	R	S	$(P \vee Q) \Rightarrow (S \vee R)$	$[(Q \Rightarrow S) \wedge (Q \Rightarrow R)] \Rightarrow [(P \vee Q) \Rightarrow (S \vee R)]$
T	T	T	T	T	T
F	T	T	T	T	T
T	F	T	T	T	T
F	F	T	T	T	T
T	T	F	T	T	T
F	T	F	T	T	T
T	F	F	T	T	T
F	F	F	T	T	T
T	T	T	F	T	T
F	T	T	F	T	T
T	F	T	F	T	T
F	F	T	F	T	T
T	T	F	F	F	T
F	T	F	F	F	T
T	F	F	F	F	F
F	F	F	F	T	T

8. (a)

P	Q	$\sim P$	$P \Rightarrow Q$	$(\sim P) \vee Q$
T	T	F	T	T
F	T	T	T	T
T	F	F	F	F
F	F	T	T	T

Since the fourth and fifth columns are the same, the propositions $P \Rightarrow Q$ and $(\sim P) \vee Q$ are equivalent.

(b)

P	Q	$P \Rightarrow Q$	$Q \Rightarrow P$	$P \Leftrightarrow Q$	$(P \Rightarrow Q) \wedge (Q \Rightarrow P)$
T	T	T	T	T	T
F	T	T	F	F	F
T	F	F	T	F	F
F	F	T	T	T	T

Since the fifth and sixth columns are the same, the propositions $P \Leftrightarrow Q$ and $(P \Rightarrow Q) \wedge (Q \Rightarrow P)$ are equivalent.

(c)

P	Q	$\sim Q$	$P \Rightarrow Q$	$\sim (P \Rightarrow Q)$	$P \wedge \sim Q$
T	T	F	T	F	F
F	T	F	T	F	F
T	F	T	F	T	T
F	F	T	T	F	F

Since the fifth and sixth columns are the same, the propositions $\sim (P \Rightarrow Q)$ and $P \wedge \sim Q$ are equivalent.

(d)

P	Q	$\sim P$	$\sim Q$	$P \wedge Q$	$\sim (P \wedge Q)$	$P \Rightarrow \sim Q$	$P \Rightarrow \sim Q$
T	T	F	F	T	F	F	F
F	T	T	F	F	T	T	T
T	F	F	T	F	T	T	T
F	F	T	T	F	T	T	T

Since the sixth, seventh and eighth columns are the same, all three propositions are equivalent.

(e)

P	Q	R	$Q \Rightarrow R$	$P \Rightarrow (Q \Rightarrow R)$	$P \wedge Q$	$(P \wedge Q) \Rightarrow R$
T	T	T	T	T	T	T
F	T	T	T	T	F	T
T	F	T	T	T	F	T
F	F	T	T	T	F	T
T	T	F	F	F	T	F
F	T	F	F	T	F	T
T	F	F	T	T	F	T
F	F	F	T	T	F	T

Since the fifth and seventh columns are the same, the propositions are equivalent.

(f)

P	Q	R	$Q \wedge R$	$P \Rightarrow (Q \wedge R)$	$P \Rightarrow Q$	$P \Rightarrow R$	$(P \Rightarrow Q) \wedge (P \Rightarrow R)$
T	T	T	T	T	T	T	T
F	T	T	T	T	T	T	T
T	F	T	F	F	F	T	F
F	F	T	F	T	T	T	T
T	T	F	F	F	T	F	F
F	T	F	F	T	T	T	T
T	F	F	F	F	F	F	F
F	F	F	F	T	T	T	T

Since the fifth and eighth columns are the same, the propositions are equivalent.

(g)

P	Q	R	$P \vee Q$	$(P \vee Q) \Rightarrow R$	$P \Rightarrow R$	$Q \Rightarrow R$	$(P \Rightarrow R) \wedge (Q \Rightarrow R)$
T	T	T	T	T	T	T	T
F	T	T	T	T	T	T	T
T	F	T	T	T	T	T	T
F	F	T	F	T	T	T	T
T	T	F	T	F	F	F	F
F	T	F	T	F	T	F	F
T	F	F	T	F	F	T	F
F	F	F	F	T	T	T	T

Since the fifth and seventh columns are the same, the propositions are equivalent.

9. (a) yes (b) no (c) yes
 (d) yes (e) no (f) no
10. (a) $[(f \text{ has a relative minimum at } x_0) \wedge (f \text{ is differentiable at } x_0)] \Rightarrow (f'(x_0) = 0)$
 (b) $(n \text{ is prime}) \Rightarrow [(n = 2) \vee (n \text{ is odd})]$
 (c) $(x \text{ is irrational}) \Rightarrow [(x \text{ is real}) \wedge \sim (x \text{ is rational})]$
 (d) $[(x = 1) \vee (x = -1)] \Rightarrow (|x| = 1)$
 (e) $(x_0 \text{ is a critical point for } f) \Leftrightarrow [(f'(x_0) = 0) \vee (f'(x_0) \text{ does not exist})]$
 (f) $(S \text{ is compact}) \Leftrightarrow [(S \text{ is closed}) \wedge (S \text{ is bounded})]$
 (g) $(B \text{ is invertible}) \Leftrightarrow (\det B \neq 0)$

- (h) $(6 \geq n - 3) \Rightarrow (n > 4) \vee (n > 10)$
- (i) $(x \text{ is Cauchy}) \Rightarrow (x \text{ is convergent})$
- (j) $(\lim_{x \rightarrow x_0} f(x) = f(x_0)) \Rightarrow (f \text{ is continuous at } x_0)$
- (k) $[(f \text{ is differentiable at } x_0) \wedge (f \text{ is strictly increasing at } x_0)] \Rightarrow (f'(x_0))$

11. (a) Let S be “I go to the store” and R be “It rains.” The preferred translation: is $\sim S \Rightarrow R$ (or, equivalently, $\sim R \Rightarrow S$). This could be read as “If it doesn’t rain, then I go to the store.”

The speaker might mean “I go to the store if and only if it doesn’t rain ($S \Rightarrow \sim R$) or possibly “If it rains, then I don’t go to the store” ($R \Rightarrow \sim S$).

- (b) There are three nonequivalent ways to translate the sentence, using the symbols D : “The Dolphins make the playoffs” and B : “The Bears win all the rest of their games.” The first translation is preferred, but the speaker may have intended any of the three.
- $\sim B \Rightarrow \sim D$ or, equivalently, $D \Rightarrow B$
 - $\sim D \Rightarrow \sim B$ or, equivalently, $B \Rightarrow D$
 - $\sim B \Leftrightarrow \sim D$ or, equivalently, $B \Leftrightarrow D$

- (c) Let G be “You can go to the game” and H be “You do your homework first.”

It is most likely that a student and parent both interpret this statement as a biconditional, $G \Leftrightarrow H$.

- (d) Let W be “You win the lottery” and T be “You buy a ticket.” Of the three common interpretations for the word “unless,” only the form $\sim T \Rightarrow \sim W$ (or, equivalently, $W \Rightarrow T$) makes sense here.

12. (a)

P	Q	R	$P \vee Q$	$(P \vee Q) \Rightarrow R$	$\sim P \wedge \sim Q$	$\sim R \Rightarrow (\sim P \wedge \sim Q)$
T	T	T	T	T	F	T
F	T	T	T	T	F	T
T	F	T	T	T	F	T
F	F	T	F	T	T	T
T	T	F	T	F	F	F
F	T	F	T	F	F	F
T	F	F	T	F	F	F
F	F	F	F	T	T	T

Since the fifth and seventh columns are the same, $(P \vee Q) \Rightarrow R$ and $\sim R \Rightarrow (\sim P \wedge \sim Q)$ are equivalent.

(b)

P	Q	R	$P \wedge Q$	$(P \wedge Q) \Rightarrow R$	$\sim Q$	$\sim R$	$P \wedge \sim R$	$(P \wedge \sim R) \Rightarrow \sim Q$
T	T	T	T	T	F	F	F	T
F	T	T	F	T	F	F	F	T
T	F	T	F	T	T	F	F	T
F	F	T	F	T	T	F	F	T
T	T	F	T	F	F	T	T	F
F	T	F	F	T	F	T	F	T
T	F	F	F	T	T	T	T	T
F	F	F	F	T	T	T	F	T

Since the fifth and ninth columns are the same, the propositions $(P \wedge Q) \Rightarrow R$ and $(P \wedge \sim R) \Rightarrow \sim Q$ are equivalent.

(c)

P	Q	R	$Q \wedge R$	$P \Rightarrow (Q \wedge R)$	$\sim Q \vee \sim R$	$(\sim Q \vee \sim R) \Rightarrow \sim P$
T	T	T	T	T	F	T
F	T	T	T	T	F	T
T	F	T	F	F	T	F
F	F	T	F	T	T	T
T	T	F	F	F	T	F
F	T	F	F	T	T	T
T	F	F	F	F	T	F
F	F	F	F	T	T	T

Since the fifth and seventh columns are the same, the propositions $P \Rightarrow (Q \wedge R)$ and $(\sim Q \vee \sim R) \Rightarrow \sim P$ are equivalent.

(d)

P	Q	R	$Q \vee R$	$P \Rightarrow (Q \vee R)$	$P \wedge \sim R$	$(P \wedge \sim R) \Rightarrow Q$
T	T	T	T	T	F	T
F	T	T	T	T	F	T
T	F	T	T	T	F	T
F	F	T	T	T	F	T
T	T	F	T	T	T	T
F	T	F	T	T	F	T
T	F	F	F	F	T	F
F	F	F	F	T	F	T

Since the fifth and seventh columns are the same, the propositions $P \Rightarrow (Q \vee R)$ and $(P \wedge \sim R) \Rightarrow Q$ are equivalent.

(e)

P	Q	R	$P \Rightarrow Q$	$(P \Rightarrow Q) \Rightarrow R$	$P \wedge \sim Q$	$(P \wedge \sim Q) \vee R$
T	T	T	T	T	F	T
F	T	T	T	T	F	T
T	F	T	F	T	T	T
F	F	T	T	T	F	T
T	T	F	T	F	F	F
F	T	F	T	F	F	F
T	F	F	F	T	T	T
F	F	F	T	F	F	F

Since the fifth and seventh columns are the same, the propositions $(P \Rightarrow Q) \Rightarrow R$ and $(P \wedge \sim Q) \vee R$ are equivalent.

(f)

P	Q	$P \Leftrightarrow Q$	$\sim P \vee Q$	$\sim Q \vee P$	$(\sim P \vee Q) \wedge (\sim Q \vee P)$
T	T	T	T	T	T
F	T	F	T	F	F
T	F	F	F	T	F
F	F	T	T	T	T

Since the third and sixth columns are the same, the propositions $P \Leftrightarrow Q$ and $(\sim P \vee Q) \wedge (\sim Q \vee P)$ are equivalent.

13. (a) If 6 is an even integer, then 7 is an odd integer.
 (b) If 6 is an odd integer, then 7 is an odd integer.
 (c) This is not possible.
 (d) If 6 is an even integer, then 7 is an even integer. (Any true conditional statement will work here.)
14. (a) If 7 is an odd integer, then 6 is an odd integer.
 (b) This is not possible.

- (c) This is not possible.
- (d) If 7 is an odd integer, then 6 is an odd integer. (Any false conditional statement will work here.)
15. (a) Converse: If $f'(x_0) = 0$, then f has a relative minimum at x_0 and is differentiable at x_0 . False: $f(x) = x^3$ has first derivative 0 but no minimum at $x_0 = 0$.
 Contrapositive: If $f'(x_0) \neq 0$, then f either has no relative minimum at x_0 or is not differentiable at x_0 . True.
- (b) Converse: If $n = 2$ or n is odd, then n is prime. False: 9 is odd but not prime.
 Contrapositive: If n is even and not equal to 2, then n is not prime. True.
- (c) Converse: If x is irrational, then x is real and not rational. True
 Contrapositive: If x is not irrational, then x is not real or x is rational. True
- (d) Converse: If $|x| = 1$, then $x = 1$ or $x = -1$. True.
 Contrapositive: If $|x| \neq 1$, then $x \neq 1$ and $x \neq -1$. True.
16. (a) tautology (b) tautology (c) contradiction
 (d) neither (e) tautology (f) neither
 (g) contradiction (h) tautology (i) contradiction
 (j) neither (k) tautology (l) neither

17. (a)

P	Q	$P \Rightarrow Q$	$\sim P$	$\sim Q$	$\sim P \Rightarrow \sim Q$
T	T	T	F	F	T
F	T	T	T	F	F
T	F	F	F	T	T
F	F	T	T	T	T

Comparison of the third and sixth columns of the truth table shows that $P \Rightarrow Q$ and $\sim P \Rightarrow \sim Q$ are not equivalent.

- (b) We see from the truth table in part (a) that both propositions $P \Rightarrow Q$ and $\sim P \Rightarrow \sim Q$ are true only when P and Q have the same truth value.
- (c) The converse of $P \Rightarrow Q$ is $Q \Rightarrow P$. The contrapositive of the inverse of $P \Rightarrow Q$ is $\sim \sim Q \Rightarrow \sim \sim P$, so the converse and the contrapositive of the inverse are equivalent.
 The inverse of the contrapositive of $P \Rightarrow Q$ is also $\sim \sim Q \Rightarrow \sim \sim P$, so it too is equivalent to the converse.

1.3 Quantifiers

1. (a) $\sim (\forall x)(x \text{ is precious} \Rightarrow x \text{ is beautiful})$ or $(\exists x)(x \text{ is precious and } x \text{ is not beautiful})$
- (b) $(\forall x)(x \text{ is precious} \Rightarrow x \text{ is not beautiful})$
- (c) $(\exists x)(x \text{ is isosceles} \wedge x \text{ is a right triangle})$
- (d) $(\forall x)(x \text{ is a right triangle} \Rightarrow x \text{ is not isosceles})$ or $\sim (\exists x)(x \text{ is a right triangle} \wedge x \text{ is isosceles})$
- (e) $(\forall x)(x \text{ is honest}) \vee \sim (\exists x)(x \text{ is honest})$
- (f) $(\exists x)(x \text{ is honest}) \wedge (\exists x)(x \text{ is not honest})$
- (g) $(\forall x)(x \neq 0 \Rightarrow (x > 0 \vee x < 0))$
- (h) $(\forall x)(x \text{ is an integer} \Rightarrow (x > -4 \vee x < 6))$ or $(\forall x \in \mathbf{Z})(x > -4 \vee x < 6)$