ATI RN MATERNAL NEWBORN PROCTORED 2023 RETAKE EXAM WITH NGN VERSION 3 QNS & ANS (VERIFIED REVISED FULL EXAM)

1. A nurse is assessing newborn following forceps assisted birth. Which of the following clinical

manifestations should the nurse identify as a complication of the birth method?

- A. Hypoglycemia
- B. Polycythemia
- C. Facial Palsy
- D. Bronchopulmonary dysplasia ANSWER-C

Which of the following statements by client indicates an understanding of the teaching?

- A. "The medication could cause me to experience heart palpitations"
- B. "This medication could cause me to experience blurred vision"
- C. "This medication could cause me to experience ringing in my ears"
- D. "This medication could cause me to experience frequent ..." ANSWER-A

A nurse is caring for a client who has hyperemesis gravidarum. Which of the following

laboratory tests should the nurse anticipate?

- A. Urine Ketones
- B. Rapid plasma regains
- C. Prothrombin time
- D. Urine culture ANSWER-A

A nurse is caring for a client who is in labor and requests nonpharmacological pain management. Which of the following nursing actions promotes client comfort? (SATA)

- A. Assisting the client into squatting position
- B. Having the client lie in a supine position
- C. Applying fundal pressure during contractions
- D. Encouraging the client to void every 6 hr ANSWER-A D

A nurse caring for a client who is at 20 weeks of gestation and has trichomoniasis. Which of

the following findings should the nurse expect?

- A. Thick, White Vaginal Discharge
- B. Urinary Frequency
- C. Vulva Lesions

D. Malodorous Discharge - ANSWER-D

A nurse is caring for a client who is 14 weeks of gestation. At which the following locations

should the nurse place the Doppler device when assessing the fetal heart rate? A. Midline 2 to 3 cm (0.8 to 1.2 in) above the symphysis pubis

- Left Upper Abdomen
- Two fingerbreadths above the umbilicus
- · Lateral at the Xiphoid Process ANSWER-A

A nurse is assessing a client who is at 27 weeks of gestation and has preeclampsia. Which of the following findings should the nurse report to the provider?

A. Urine protein contraction 200mg/24 hours

- B. Creatine 0.8
- C. Hemoglobin 14.8
- D. Platelet 60,000 ANSWER-D

A nurse is teaching about clomiphene citrate to a client who is experiencing infertility. Which of the following adverse effects should the nurse include?

- Tinnitus
- Urinary Frequency
- C. Breast Tenderness
- D. Chills ANSWER-C (vision disturbances).

A nurse is assessing a newborn upon admission to the nursery. Which of the following should the nurse expect?

A. Bulging fontanelles

- B. Nasal flaring
- C. Length from head to heel of 40cm
- D. Chest circumference 2c (smaller than the head circumference ANSWER-D

A nurse is planning care for a newborn who has necrotizing abstinence syndrome. Which of the following interventions should the nurse include in the plan of care? A.Increase the newborn visual stimulation

- B. Weigh the newborn everyday
- C. Discourage parental interaction after a social evaluation
- D. Swaddle the newborn in a flexed position ANSWER-D

A nurse is caring for a newborn who his 6 hours old had has a bedside glucometer reading of 65mg/dl. The newborns mother has type 2 diabetes mellitus. Which of the following actions should the nurse take.

A. obtain a blood sampling for a serum glucose level.

- B. Feed the newborn immediately
- C. Administer 50ml of dextrose
- D. Reassess the blood glucose reading prior to the next feeding ANSWER-D

A nurse is providing teaching to a client about exercise safety during pregnancy. Which of the folioing statements by the client indicates an understanding of the teaching (SATA)?

- I will limit my time in the hot tub to 30 miniutes after exercise
- · I should consume three 8 ounce glasses of water after I exercise
- C. I will check my heart rate every 15 minutes during
- D. I should limit exercise sessions to 30 minutes when the weather is humid

E. I should rest by lying on my side for 10 minutes following exercise. - ANSWER-B D E

A charge nurse is teaching a group of staff nurses about fetal monitoring during labor. Which of the following findings should the charge nurse instruct the staff members to report to the provider?

- A. Contractions duration 95 seconds to 100 seconds
- B. Contraction frequency 2-3 minutes
- C. Absent early decelerations of fetal heart rate ANSWER-C

A nurse in a woman's health clinic is obtaining a health history from a client. Which of the following findings should the nurse identify as increasing the client's risk for developing pelvic inflammatory disease (PID)?

A. Recurrent Cystitis

- Frequent Alcohol Use
- Use of Oral Contraceptives
- Chlamydia Infection ANSWER-D

A nurse is teaching a prenatal class about immunizations that newborns receive following

birth. Which of the following immunizations should the nurse include in the teaching?

- A. Hepatitis B
- Rotavirus
- Pneumococcal
- Varicella ANSWER-A

A nurse is providing nutritional guidance to a client who is pregnant and follows a vegan

diet. The client asks the nurse which foods she should eat to ensure adequate calcium intake. The

nurse should instruct the client that which of the following foods has the highest amount of

calcium?

A. ¹/₂ cup cubed avocado

- B. 1 large banana
- C. 1 medium potato
- D. 1 cup cooked broccoli ANSWER-D

A nurse in a provider's office is assessing a client at her first antepartum visit. Theclient states that the first day of her last menstrual period was March 8. - ANSWER-December 15

A nurse is caring for a client who is in the second stage of labor. Which of the following

manifestations should the nurse expect?

- A. The client expels the placenta
- B. The client experiences gradual dilation of the cervix
- C. The client begins have regular contractions.
- D. The client delivers the newborn ANSWER-D

A nurse is assessing a client who is at 37 weeks of gestation. Which of the following statements by the client requires immediate intervention by the nurse?

- "It burns when I urinate
- "My feet are really swollen today". ANSWER-A

A nurse is providing discharge teaching to a new parent about car seat safety. Which of the

following statements by the parent indicates an understanding of the teaching?

- A. "I should position my baby's car seat at a 45 degree angle in the car."
- B. "I should place the car seat rear facing until my baby is 12 months old."
- C. "I should place the harness snugly in a slot above my baby's sholders."
- D. "I should position the retainer clip at the top of my baby's abdomen. ANSWER-B

A nurse on an antepartum unit is reviewing the medical records for four clients. Which of the following clients should the nurse assess first?

- A client who has diabetes mellitus and an HbA1c of 5.8%
- A client who has preeclampsia and a creatinine level of 1.1 mg/ dL
- · A client who has hyperemesis gravidarum and a sodium level of 110 mEq/L

D.

• A client who has placenta previa and a hematocrit of 36% - ANSWER-C

A nurse on the postpartum unit is caring for four clients. For which of the following clients

should the nurse notify the provider?

- A client who has a urinary output of 300 ml in 8 hr
- · A client who reports abdominal cramping during breastfeeding
- A client who is receiving magnesium sulfate and has absent deep tendon reflexes

 A client who reports lochia rhubra requiring changing perineal pads every 3 h -ANSWER-C

A nurse is caring for a client who has active genital herpes simplex virus type 2. Which of the following medications should the nurse plan to administer?

- A. Metronidazole
- B. Penicillin
- C. Acyclovir
- D. Gentamicin ANSWER-C

A nurse is caring for a client following an amniocentesis. The nurse should observe the client for which of the

following complications?

- A. Hyperemesis
- B. Proteinuria
- C. Hypoxia
- D. Hemorrhage ANSWER-D

A nurse is planning care for a client who is receiving oxytocin by continuous IV infusion for labor induction.

Which of the following interventions should the nurse include in the plan?

A. Increase the infusion rate every 30 to 60 min.

- B. Maintain the client in a supine position.
- C. Titrate the infusion rate by 4 milliunits/min.
- D. Limit IV intake to 4 L per 24 hr ANSWER-A