

NURS 6635 Midterm Exam NO Answer

Nursing Education (Walden University)

NURS 6635/6640 Midterm Exam

QUESTION 1

1. Select the theorist who is credited with establishing the area of psychosomatic medicine.

- ~~A.~~ Freud
- B. Abraham
- C. Groddeck
- D. Ferenczi**

1 point

QUESTION 2

1. Select the most important component of the patient-provider rapport.

- A. Empathy**
- B. Confidentiality
- C. Respect
- D. Privacy

1 point

QUESTION 3

1. Select the disorder in which patients feel detached from their environment.

- ~~A.~~ Derealization
- B. Transient global amnesia

- C. Depersonalization
- D. Dissociative identity

1 point

QUESTION 4

1. Select the patient evaluation scale to be used before and during prescribing antipsychotic drugs.

- A. MMSE
- B. AIMS
- C. CAGE
- D. HAM-D

1 point

QUESTION 5

1. Select the event most associated with the occurrence of dissociative fugue.

- A. Head trauma
- B. Substance abuse
- C. Partner betrayal
- D. Unplanned travel

1 point

QUESTION 6

1. Select the most influential source of bias in a mental status examination.

- A. Culture

- B. Genetics
- C. Environment
- D. Age

1 point

QUESTION 7

1. Select the nondrug therapy with the greatest evidence of effectiveness in treating major depressive disorder.
- ~~A.~~ Family
 - B. Psychoanalytic
 - C. Cognitive**
 - D. Behavior

1 point

QUESTION 8

1. Select the therapy that has been most effective in treating childhood PTSD.
- ~~A.~~ SSRI antidepressants
 - B. CBITS**
 - C. SPARCS
 - D. Crisis intervention

1 point

QUESTION 9

1. Select the rating scale used to assess response to antidepressant drug therapy.

- A. BDI
- B. MMSE
- C. BPRS
- D. HAM-D**

1 point

QUESTION 10

1. Select the greatest impediment to treating anorexia nervosa patients.
- ~~A.~~ Drug adverse effects
 - B. Variability of family therapy
 - C. Patient resistance**
 - D. Noncompliance with therapy

1 point

QUESTION 11

1. Select the public performance that is most associated with social anxiety in children.
- ~~A.~~ Dancing
 - B. Speaking**
 - C. Athletics
 - D. Eating

1 point

QUESTION 12

1. Select the difference in rates of suicidal thoughts and behaviors between placebo and drug in children and adolescents observed from random controlled trials of nine antidepressant drugs that resulted in the 2004 FDA “black box” warning in labeling of antidepressant drugs.
 - A. 1% placebo, 5% drug
 - B. 2% placebo, 4% drug
 - C. 2% placebo, 10% drug
 - D. 4% placebo, 12% drug

1 point

QUESTION 13

1. Select the drug used to treat bipolar disorder that may cause hypothyroidism.
 - A. Lithium
 - B. Risperidone
 - C. Lamotrigine
 - D. Valproate

1 point

QUESTION 14

1. Select the rating scale area developed by the World Health Organization (WHO).
 - A. Cognitive
 - B. Anxiety
 - C. Depression

D. Disability

1 point

QUESTION 15

1. Select the rating scale used to screen for Alzheimer's disease and other dementias.

- A. BPRS
- B. YBOCS
- C. NPI
- D. BDI

1 point

QUESTION 16

1. Select the mood disorder that occurs most frequently in children and adolescents with ADHD.

- A. Bipolar
- B. Major depression
- C. Dysthymia
- D. Conduct

1 point

QUESTION 17

1. Select the activity in which children are most likely to suffer selective mutism.

- A. Religious

- B. Family
- C. Playmates
- D. School**

1 point

QUESTION 18

1. Select the comorbidity most commonly occurring with intermittent explosive disorder.

- ~~A.~~ Narcissism
- B. Pyromania
- C. Depersonalization
- D. Kleptomania

1 point

QUESTION 19

1. Select the quantities of mental disorders categories and discrete illnesses in DSM-5.

- ~~A.~~ 10 and 60
- B. 16 and 120
- C. 22 and 150
- D. 31 and 175

1 point

QUESTION 20

1. Select the scientist who is most associated with the development of learning theory.

- A. Piaget
- B. Ainsworth
- C. Bowlby
- D. Pavlov

1 point

QUESTION 21

1. Select the short rating scale used to assess change in psychiatric inpatients.

- A. PANSS
- B. BPRS
- C. HAM-D
- D. CAPS

1 point

QUESTION 22

1. Select the period of time after most recent use that THC (marijuana) can be detected in urine specimens.

- A. 7-12 hours
- B. 36-72 hours
- C. 3 days
- D. 2-7 days

1 point

QUESTION 23

1. Select the therapy for insomnia that can be effective at up to three years after discontinuation.

- A. Cognitive-behavioral
- B. Sedating antihistamines
- C. Benzodiazepines
- D. Paradoxical intention

1 point

QUESTION 24

1. Select **two** brain regions that are associated with anxiety disorders.

- A. Substantia nigra
- B. Locus ceruleus
- C. Temporal lobe
- D. Raphe nuclei

1 point

QUESTION 25

1. Select the hormone or neurotransmitter that is involved in regulating the 24-hour circadian sleep-wake cycle.

- A. Dopamine
- B. Corticotrophin releasing

C. Serotonin

D. Melatonin

1 point

QUESTION 26

1. Select the medical condition that must be excluded as a cause of somatic symptom disorder or hypochondriasis.

A. Angina pectoris

B. Asthma

C. Diabetes

D. Multiple sclerosis

1 point

QUESTION 27

1. Select the **two** rating scales used specifically for schizophrenia.

A. PANSS

B. MMSE

C. SAPS

D. BPRS

1 point

QUESTION 28

1. Select **two** events that are causes of dissociative trance disorder.

2. A. Natural disaster

B. Substance abuse

C. Experienced violence

D. Appropriate drug therapy

1 point

QUESTION 29

1. Select the term that best describes the study of human behavior based on transmission and modification of genetically influenced behavioral traits.

~~A.~~ Ethology

B. Sociobiology

C. Neuropsychology

D. Phenomenology

1 point

QUESTION 30

1. Select the theorist who developed psychoanalysis.

~~A.~~ Maslow

B. Skinner

C. Erikson

D. Freud

1 point

QUESTION 31

1. Select the serum assessment that is required when prescribing drugs to treat psychiatric disorders in psychosomatic patients.

- A. Potassium
- B. White blood cells
- C. Liver enzymes
- D. Blood urea nitrogen

1 point

QUESTION 32

1. Select **two** types of theories that have contributed to causes of anxiety.

- A. Psychoanalytic theory
- B. Behavioral theory
- C. Family
- D. Group

1 point

QUESTION 33

1. Select the neurotransmitter that is most genetically involved in childhood and adolescent depression.

- A. Norepinephrine
- B. Serotonin
- C. Dopamine
- D. Gamma amino butyric acid (GABA)

1 point

QUESTION 34

1. Select the drug that has been effective in treating social anxiety in children.
- A. Bupropion
 - B. Carbamazepine
 - C. Amitriptyline
 - D. Fluoxetine

1 point

QUESTION 35

1. Select the alternate name for persistent depressive disorder.
- A. Cyclothymia
 - B. Bipolar I
 - C. Bipolar II
 - D. Dysthymia

1 point

QUESTION 36

1. Select the percentage occurrence of anxiety disorders in children and adolescents.
- A. 2-5
 - B. 5-10
 - C. 10-20

D.

20-30

1 point

QUESTION 37

1. Select the rating scale used to assess obsessive-compulsive disorder.

~~A.~~

HAM-A

B.

CAGE

C.

YBOCS

D.

CAPS

1 point

QUESTION 38

1. Select the characteristic in long sleepers (more than 9 hours) that is absent in short sleepers (less than 6 hours).

~~A.~~

Mildly depressed

B.

Ambitious

C.

Socially adept

D.

Efficient

1 point

QUESTION 39

1. Select the function that is most affected by conversion disorder.

A.

Voluntary motor

B.

Hearing

C. Involuntary motor

D. Smelling

1 point

QUESTION 40

1. Select the minimum treatment time to assess the effectiveness of antidepressant drugs

A. 1-2 weeks

B. 3-4 weeks

C. 6-8 weeks

D. 10-14 weeks

1 point

QUESTION 41

1. Select the largest percentage prevalence of psychiatric disorders in patients hospitalized for medical conditions.

2. A. 10

B. 33

C. 50

D. 67

1 point

QUESTION 42

1. Select the percent rate of mortality in anorexia nervosa patients.

- A. 1 to 5
- B. 3 to 12
- C. 5 to 18
- D. 18 to 25

1 point

QUESTION 43

1. Select the therapeutic plasma concentration range for lithium in mEq/L.

- ~~A.~~ 0.1-0.5
- B.** 0.6-1.2
- C. 1.4-1.9
- D. 2.0-2.6

1 point

QUESTION 44

1. Select the type of dissociative amnesia described by patient's inability to recall any information about her or his family.

- ~~A.~~ Selective
- B. Localized
- C. Continuous
- D. Systematized

1 point

QUESTION 45

1. Select the memory category retained in transient global amnesia that is absent in dissociative amnesia.

- A. Family members
- B. Sequential events
- C. Personal identity
- D. Adolescent years

1 point

QUESTION 46

1. Select the behavior that best describes early onset bipolar disorder in children and adolescents.

- A. Aggression
- B. Withdrawal
- C. Panic
- D. Exhibition

1 point

QUESTION 47

1. Select the serum substance that should be measured to aid the diagnosis of restless leg syndrome.

- A. Potassium
- B. Ferritin
- C. Magnesium

D. Albumin

1 point

QUESTION 48

1. Select the **two** primary psychometric properties of psychiatric rating scales.

~~A.~~ Reproducibility

B. Reliability

C. Validity

D. Variability

1 point

QUESTION 49

1. Select the third most commonly reported psychiatric symptom.

~~A.~~ Depression

B. Anxiety

C. Psychosis

D. Depersonalization

1 point

QUESTION 50

1. Select the anxiety disorder that emerges in ages 9 to 18 months.

A. Separation

B. Generalized

- C. Social
- D. Traumatic

1 point

QUESTION 51

1. Select the factor that best correlates with the occurrence of PTSD in children.
- ~~A.~~ Excessive dopamine metabolites
 - B. Higher IQ
 - C. Larger corpus callosum
 - D. Lower IQ**

1 point

QUESTION 52

1. Select the percentage that most accurately represents the rate of heritability of major depression.
2.
 - A. 15
 - B. 30
 - C. 45
 - D. 60**

1 point

QUESTION 53

1. Select the least effective method to treat dissociative amnesia.
- ~~A.~~ Antipsychotic drugs