

UPDATED HESI OB MATERNITY EXAM QUESTIONS AND ANSWERS

A client at 37 weeks gestation presents to labor and delivery with contractions every two minutes the nurse observes several shallow small vesicles on her pubis labia and perineum. the nurse should recognize the clients is prohibiting symptoms of which condition?

German measles

herpes simplex virus

syphilis

genital warts

A client who had her first baby three months ago and is breastfeeding her infant tells the nurse that she is currently using the same diaphragm that she used before becoming pregnant. Which information should the nurse provide this client?

Use alternative form of birth control until new diaphragm can be obtained.

A 30- year-old primigravida delivers a 9-pound infant vaginally after a 30-hour labor. What is the priority nursing action for this client?

Observe for signs of uterine hemorrhage

At 0600 while admitting a woman for a scheduled repeat cesarean section (C- Section), the client tells the nurse that she drank a cup a coffee at 0400 because she wanted to avoid getting a headache. Which action should the nurse take first?

Inform the anesthesia care provider

The nurse is caring for a postpartum client who is exhibiting symptoms of spinal headache 24 hours following delivery of a normal newborn. Prior to the anesthesiologist arrival on the unit, which action should the nurse

perform?

- Place procedure equipment at bedside

The nurse is caring for a newborn who is 18 inches long, weighs 4 pounds, 14 ounces, has a head circumference of 13 inches, and a chest circumference of 10 inches. Based on these physical findings, assessment for which condition has the highest priority?

Hypoglycemia

12.12.

the nurse is caring for a 35 week gestation infant delivered by cesarean section 2 hours ago. the nurse observes the infants respiratory rate is 72 breaths/minute with nasal flaring, grunting, and retractions. the nurse should recognize these findings indicate which complication?

- B – transient tachypnea of the newborn (double check this answer)

13.

A primipara client at 42 weeks gestation is admitted for induction. within one hour after initiating an oxytocin infusion, her cervix is 100% effaced and 6 cm dilated, contractions are occurring every 1 minute with a 75 second duration. when the nurse stops the oxytocin and starts oxygen. after 30 minutes of uterine rest, the contractions are occurring every 5 minutes with 20 second duration. which intervention should the nurse implement?

A primigravida arrives at the observation unit of the maternity unit because she thinks she is in labor. the nurse applies the external fetal heart monitor and determines

A primigravida client with gestational hypertension and a Bishop score of 3 is scheduled for induction of labor. the nurse administers misoprostol at 0700 then