

MULTIPLE-CHOICE TEST ITEMS

CHAPTER 1

INTRODUCTION

1.1 Statistics exists because

- a) variability.
- b) mathematics.
- c) complexity.
- d) commonalities.

Ans: a

1.2 Which of the following is *not* one of the reasons mentioned for taking an introductory statistics class?

- a) better understand research reports in your special area of interest
- b) plan statistical analyses for modest research projects
- c) intelligently evaluate statistical references in news publications and TV broadcasts
- d) single-handedly plan the analysis for any research project

Ans: d

1.3 The more advanced area of statistics is

- a) descriptive statistics.
- b) inferential statistics.
- c) population statistics.
- d) analytical statistics.

Ans: b

1.4 The area of statistics that *organizes and summarizes* information about a collection of actual observations is known as

- a) descriptive statistics.
- b) inferential statistics.
- c) population statistics.
- d) analytical statistics.

Ans: a

1.5 A single word that best describes *inferential* statistics is

- a) analyzing.
- b) summarizing.
- c) organizing.
- d) generalizing.

Ans: d

1.6 Indicate whether *one, both, or neither* of the following statements typifies *descriptive* statistics.

- a) It usually takes several months before a person feels "at home" in a new environment.
- b) My income last summer was about \$10,000.
- c) both a and b
- d) neither a nor b

Ans: b

1.7 Indicate whether *one, both, or neither* of the following statements typifies *descriptive* statistics.

- a) There is a tendency for elderly people to postpone their death until after their birthday.
- b) People with similar personalities are mutually attracted.
- c) both a and b
- d) neither a nor b

Ans: d

1.8 Indicate whether *one, both, or neither* of the following statements typifies *inferential* statistics.

- a) Daily meditation reduces stress.

- b) The national deficit for last year exceeded 500 billion.
- c) both a and b
- d) neither a nor b

Ans: a

1.9 When conducting a survey, it is important that the sample be

- a) large.
- b) haphazard
- c) carefully selected
- d) random

Ans: d

1.10 Some form of randomization should occur in both

- a) descriptive and inferential statistics
- b) populations and samples
- c) surveys and experiments
- d) independent and dependent variables

Ans: c

1.11 Random assignment helps us to determine whether an observed difference between two groups is

- a) larger than expected just by chance.
- b) probably is real.
- c) merits further attention.
- d) all of the above.

Ans: d

1.12 A statistical analysis is based on

- a) data.
- b) words.
- c) labels.
- d) numbers.

Ans: a

1.13 *Quantitative* observations consist of

- a) words.
- b) numerical codes.
- c) numbers.
- d) all of the above

Ans: c

1.14 Indicate whether *one, both, or neither* of the following statements describes *quantitative* observations.

- a) score on this exam
- b) vocational goal
- c) both a and b
- d) neither a nor b

Ans: a

1.15 Indicate whether *one, both, or neither* of the following statements describes *qualitative* observations.

- a) place of birth
- b) political preference
- c) both a and b
- d) neither a nor b

Ans: c

1.16 Indicate whether *one, both, or neither* of the following statements describes *qualitative* observations.

- a) IQ score
- b) age
- c) both a and b
- d) neither a nor b

Ans: d

1.17 Indicate whether *one, both, or neither* of the following statements describes *ranked* observations.

- a) finish order at a car race.
- b) birth order among children in a family
- c) both a and b
- d) neither a nor b

Ans: c

1.18 In a survey of religious affiliation, numbers are assigned as follows: 1-None, 2-Christian, 3-Jewish, 4-Buddhist, 5-Other. Therefore it is appropriate to conclude that

- a) two Christians equal one Buddhist.
- b) a Jew is intermediate between a Christian and a Buddhist.
- c) five different classes of religious affiliation are being distinguished.
- d) religious affiliation can be treated as quantitative data.

Ans: c

1.19 An important *first* step in a statistical analysis requires that observations be identified as either

- a) words or numerical codes.
- b) quantitative, ranked, or qualitative.
- c) true or false.
- d) amounts or counts.

Ans: b

1.20 Data are *quantitative* if any single observation within a batch of observations represents a(n)

- a) amount or count.
- b) word or label.
- c) coding device.
- d) bit of information.

Ans: a

1.21 You are asked whether 1, 3, 2, 5, 3, 7 constitute quantitative or qualitative data. Your best reply is

- a) *quantitative* because these numbers involve an amount or count.
- b) *qualitative* because these numbers reflect arbitrary numerical codes or labels.
- c) *quantitative or qualitative* depending on the accuracy of these numbers.
- d) *quantitative or qualitative* depending on whether these numbers represent an amount or count, or merely a numerical code.

Ans: d

1.22 The simplest level of measurement is

- a) interval/ratio.
- b) ordinal.
- c) nominal.
- d) approximately interval.

Ans: c

1.23 The most complex level of measurement is

- a) interval/ratio
- b) ordinal.

- c) nominal.
- d) approximately interval.

Ans: a

1.24 Shifts to more complex levels of measurement are accompanied by sets of observations that contain

- a) more information.
- b) less information.
- c) more errors.
- d) fewer errors.

Ans: a

1.25 If movies are rated on a scale from four stars (outstanding) to no stars (terrible), measurement is

- a) interval/ratio
- b) ordinal.
- c) nominal.
- d) approximately interval.

Ans: b

1.26 If people are classified as either literate, semi-literate, or illiterate, measurement is

- a) interval/ratio
- b) ordinal.
- c) nominal.
- d) approximately interval.

Ans: b

1.27 If college students are polled about how many academic units they are carrying during the current term, measurement is

- a) interval/ratio
- b) ordinal.
- c) nominal.
- d) approximately interval.

Ans: a

1.28 Qualitative data are associated with

- a) all levels of measurement.
- b) interval/ratio, ordinal, and nominal measurement.
- c) ordinal and nominal measurement.
- d) nominal measurement.

Ans: c

1.29 Quantitative data are associated with

- a) all levels of measurement.
- b) interval/ratio and ordinal measurement.
- c) interval/ratio
- d) interval/ratio and approximately interval measurement.

Ans: d

1.30 Ranked data are associated with

- a) all levels of measurement
- b) ordinal measurement
- c) approximately interval measurement
- d) nominal measurement.

Ans: b

1.31 The distinctive property of ordinal measurement is

- a) equal intervals.

- b) order.
- c) classification.
- d) a true zero.

Ans: b

1.32 Which level of measurement is *not* represented in the following statement? A racehorse, wearing number *three*, finishes *second*, with a time of *1.50 minutes*.

- a) nominal
- b) ordinal
- c) approximately interval
- d) interval/ratio

Ans: c

1.33 Which level of measurement is represented twice in the following statement? In my group dynamics class, I arrived *last* but spoke *most often* during the lengthy *three-hour* session.

- a) nominal
- b) ordinal
- c) approximately interval
- d) interval/ratio

Ans: b

1.34 Six is twice three only when these numbers emerge from a scale of measurement having

- a) classification.
- b) order.
- c) equal intervals.
- d) a true zero.

Ans: d

1.35 The attainment of interval/ratio measurement is particularly difficult when you attempt to measure

- a) physical characteristics.
- b) nonphysical characteristics.
- c) complex characteristics.
- d) simple characteristics.

Ans: b

1.36 As measures of academic achievement, grade point averages only approximate interval measurement. Nevertheless, it would be permissible to claim that a GPA of 2.00 represents

- a) an amount of academic achievement roughly midway between GPAs of 1.00 and 3.00.
- b) twice as much academic achievement as a GPA of 1.00.
- c) an amount of academic achievement midway between GPAs of 1.00 and 3.00.
- d) none of the above

Ans: a

1.37 Data that approximate interval measurement receive the same statistical treatment as

- a) nominal and ordinal data.
- b) ordinal and interval data.
- c) interval/ratio data.
- d) ordinal, interval, and interval/ratio data.

Ans: c

1.38 When data only approximate interval measurement, as often happens in the behavioral and social sciences, you should

- a) interpret numerical claims cautiously.
- b) shift to more precise measurement.
- c) question the worth of the data.
- d) develop more incisive research techniques

Ans: a

1.39 A characteristic that can assume more than one value is referred to as

- a) fickle.
- b) changeable.
- c) a constant.
- d) a variable.

Ans: d

1.40 Which one of the following quantitative variables is *not* continuous?

- a) age
- b) speed
- c) population
- d) height

Ans: c

1.41 When values are rounded off, the resulting numbers are

- a) approximate.
- b) erroneous.
- c) misleading.
- d) speculative.

Ans: a

1.42 Gaps among values of continuous variables are

- a) more apparent than real.
- b) caused by rounding off procedures.
- c) reflect our need to deal with finite numbers.
- d) all of the above.

Ans: d

1.43 To determine whether a new sleeping pill is effective, adult insomniacs receive a pill (either real or fake, according to some impartial assignment rule) and subsequently their sleeping times are measured, in minutes, during eight-hour observation periods. In this study, sleeping time is

- a) the independent variable.
- b) the dependent variable.
- c) either the independent or the dependent variable.
- d) neither the independent nor the dependent variable.

Ans: b

1.44 To determine whether a new sleeping pill is effective, adult insomniacs receive a pill (either real or fake, according to some impartial assignment rule) and subsequently their sleeping times are measured, in minutes, during eight-hour observation periods.

This study can *best* be described as

- a) an experiment.
- b) an observational study.
- c) one involving two variables.
- d) one involving human subjects.

Ans: a

1.45 An independent variable is defined as a treatment that the investigator

- a) measures.
- b) manipulates.
- c) modifies.
- d) makes.

Ans: b

1.46 A distinctive property of an experiment is that the investigator decides on

- a) the laboratory setting.

- b) the two variables to be studied.
- c) the quantification of the dependent variable.
- d) who receives the special treatment.

Ans: d

1.47 When compared to observational studies, well-designed experiments provide conclusions that are more clear-cut about

- a) human populations.
- b) relationships.
- c) cause-effect relationships.
- d) large batches of data.

Ans: c

1.48 When variables cannot be manipulated by the investigator, relationships must be studied with

- a) observational studies
- b) patience
- c) very small numbers of subjects
- d) an abstract perspective

Ans: a

1.49 A confounding variable

- a) increases the generality of a study
- b) compromises the interpretation of a study
- c) replaces the independent variable
- d) facilitates the interpretation of a study

Ans: b

1.50 An experiment permits a decision about whether an observed difference is

- a) true or false.
- b) large or small.
- c) real or transitory.
- d) important or unimportant.
- e) Ans: c

MULTIPLE-CHOICE TEST ITEMS

CHAPTER 2

DESCRIBING DATA WITH TABLES AND GRAPHS

2.1 In a frequency distribution for *ungrouped* data,

- a) class intervals are of size one.
- b) observations are oversummarized.
- c) duplicate observations can't occur.
- d) the range of possible observations should be very large.

Ans: a

2.2 Frequency distributions can be constructed for

- a) grouped data.
- b) ungrouped data.
- c) qualitative data.

d) all of the above

Ans: d

2.3 To organize a set of observations, such as 2, 3, 0, 5, 8, 1, 3, 1, for number of children per household, you should use a frequency distribution for *ungrouped* data because

- a) the number of possible values is small.
- b) observations are whole numbers.
- c) observations tend to cluster together.
- d) data are quantitative.

Ans: a

2.4 In a frequency distribution for grouped data, the leftmost column consists of

- a) frequencies.
- b) all possible values.
- c) all observed values.
- d) class intervals.

Ans: d

2.5 One inevitable byproduct of grouping data is the loss of

- a) the identities of individual observations.
- b) some data.
- c) some regularities or patterns in the data.
- d) all of the above

Ans: a

2.6 Identify the one *optional* guideline for a well-constructed frequency distribution.

- a) All observations should be included in one, and only one, class.
- b) All classes (with both upper and lower boundaries) should be equal in width.
- c) All classes--even those with zero frequencies--should be listed.
- d) All classes should have both boundaries.

Ans: d

2.7 Which one of the following *is not* a guideline for a well-constructed frequency distribution?

- a) Select the width of the class from convenient numbers.
- b) Aim for approximately ten classes.
- c) The lower boundary of the bottom class should coincide with the smallest observation.
- d) All classes should have both boundaries.

Ans: c

2.8 What's wrong with the following frequency distribution?

\underline{X}	\underline{f}
200-249	23
150-199	32
100-149	16
50- 99	<u>7</u>
	78

- a) unequal classes
- b) gaps between classes
- c) exclusion of zero
- d) too few classes

Ans: d

2.9 What's wrong with the following frequency distribution?

\underline{X}	\underline{f}
220-239	1
200-219	0
160-199	0

140-159	11
120-139	17
100-119	21
70- 99	14
60- 69	8
30- 59	7
20- 29	2
0- 19	<u>1</u>
	82

- a) gaps between classes
- b) unequal classes
- c) not exactly ten classes
- d) nothing is wrong

Ans: b

2.10 Too little summarization tends to occur if the frequency distribution contains

- a) too many classes.
- b) too few classes.
- c) unequal classes.
- d) open-ended classes.

Ans: a

2.11 In a well-constructed frequency distribution, the size of the gap between the boundaries of classes always equals

- a) the width of the class.
- b) the unit of measurement.
- c) some convenient number.
- d) a value of one.

Ans: b

2.12 To determine the real limits of a class interval,

- a) ignore gaps between intervals.
- b) simply find the difference between tabled boundaries.
- c) add the tabled boundaries and divide by two.
- d) locate the midpoints in gaps between tabled boundaries.

Ans: d

2.13 In the following well-constructed frequency distribution,

<u>X</u>	<u>f</u>
135-149.99	1
120-134.99	0
105-119.99	1
90-104.99	1
75- 89.99	2
60- 74.99	<u>5</u>
	10

the unit of measurement equals

- a) 0.01
- b) 1.00
- c) 13.99
- d) 15.00

Ans: a

2.14 (NOTE: This question requires display of frequency distribution in 2.13.)

The class interval width equals

- a) 13.99
- b) 15
- c) 60
- d) none of the above

Ans: b

2.15 (NOTE: This question requires display of frequency distribution in 2.13.)

The overall appearance or shape of this distribution is

- a) balanced.
- b) lopsided.
- c) regular.
- d) impossible to determine from the information given.

Ans: b

2.16 (NOTE: This question requires display of frequency distribution in 2.13.)

The relative frequency of the interval 75-89.99 equals

- a) 0.10
- b) 0.20
- c) 0.50
- d) none of the above

Ans: b

2.17 (NOTE: This question requires display of frequency distribution in 2.13.)

The approximate percentile rank for the interval 90-104.99 is

- a) 10
- b) 70
- c) 80
- d) none of the above

Ans: c

2.18 (NOTE: This question requires display of frequency distribution in 2.13.)

If this frequency distribution represents the test scores for a class of ten college students, we can infer that the test probably is

- a) too easy.
- b) too difficult.
- c) multiple choice.
- d) essay.

Ans: b

2.19 (NOTE: This question requires display of frequency distribution in 2.13.)

The cumulative frequency for the interval 105-119.99 equals

- a) 1
- b) 9
- c) 10
- d) none of the above

Ans: b

2.20 Among a batch of 95 observations, the smallest observation equals 300 and the largest observation equals 349. Therefore, the class interval width should equal

- a) 1
- b) 3
- c) 5
- d) 10

Ans: c

2.21 If the smallest observation equals 743 and the class interval width is to equal 50, the boundaries of the bottom class should be

- a) 700-749
- b) 740-789
- c) 743-792
- d) 750-799

Ans: a

2.22 Outliers are

- a) worthless observations.
- b) inaccurate observations.
- c) illegal observations.
- d) very extreme observations.

Ans: d

2.23 Whenever an outlier is encountered, *first* attempt to

- a) verify its accuracy.
- b) segregate it from any summary of data.
- c) relegate it to a footnote.
- d) study the special circumstances that produce it.

Ans: a

2.24 A valid outlier

- a) is to be avoided at all costs.
- b) might enhance your understanding.
- c) occurs in almost every batch of data.
- d) indicates poor data-collecting techniques.

Ans: b

2.25 Relative frequency distributions allow us to focus on

- a) the part or fraction of the total frequency that occupies each class.
- b) the relative density of observations among difference classes or categories within the same distribution.
- c) the shapes of two or more distributions based on different total numbers of observations.
- d) all of the above

Ans: d

2.26 To convert a frequency distribution into a relative frequency distribution,

- a) add the frequencies of all lower intervals.
- b) change proportions to percents.
- c) divide the frequency for each interval by the total frequency for the entire distribution.
- d) move the decimal point two places to the right.

Ans: c

2.27 Cumulative frequencies indicate

- a) how many observations fall at or below a particular class.
- b) how many observations fall at or above a particular class.
- c) either a or b depending on your perspective.
- d) neither a nor b

Ans: a

2.28 When relative standing within a distribution assumes primary importance, as with some test scores, frequencies often are converted to

- a) relative frequencies.
- b) cumulative frequencies.
- c) cumulative relative frequencies.
- d) all of the above

Ans: c

2.29 The percentile rank of an observation refers to its

- a) rank in the distribution.
- b) cumulative proportion in the distribution.
- c) cumulative percent in the distribution.
- d) cumulative frequency in the distribution.

Ans: c

2.30 For a test of math achievement, it would be preferable to attain a percentile rank of

- a) 1
- b) 43
- c) 54
- d) 76

Ans: d

2.31 An androgynous person (that is, a person without either pronounced male or female tendencies) should have a percentile rank in the vicinity of _____ on a masculinity-femininity scale.

- a) 10
- b) 30
- c) 50
- d) 70

Ans: c

2.32 The assignment of *exact* percentile ranks requires

- a) excessive computations.
- b) ungrouped data.
- c) accurate computations.
- d) a large total number of observations.

Ans: b

2.33 In the following frequency distribution,

<u>X</u>	<u>f</u>
6	1
5	2
4	3
3	0
2	0
1	<u>3</u>
	10

an X value of 1 has a percentile rank of

- a) 4
- b) 30
- c) 40
- d) 60

Ans: b

2.34 (NOTE: This question requires display of the frequency distribution in 2.33.) In the frequency distribution, a percentile rank of 60 is paired with an X value of

- a) 3
- b) 3.5
- c) 4
- d) 5

Ans: c

2.35 For the following frequency distribution,

<u>X</u>	<u>f</u>
80-89	4
70-79	2
60-69	11
50-59	<u>3</u>
	20

the interval 70-79 has an approximate percentile rank of

- a) 20
- b) 30
- c) 80
- d) none of the above

Ans: c

2.36 Frequency distributions for qualitative data are relatively easy to construct because

- a) classes or categories are dictated by the data.
- b) they usually are based on small total numbers of observations.
- c) data patterns are more detectable.
- d) all of the above

Ans: a

2.37 Frequency distributions for qualitative data always can be converted into

- a) relative frequency distributions.
- b) cumulative frequency distributions.
- c) cumulative relative frequency distributions.
- d) all of the above

Ans: a

2.38 It wouldn't make sense to determine _____ for the frequency distribution of ethnic groups within the state of New York.

- a) cumulative frequencies
- b) cumulative relative frequencies
- c) percentile ranks
- d) any of the above

Ans: d

2.39 Which one of the following *isn't* a frequency distribution for qualitative data?

- a) type of transportation used by college students
- b) length of marriages at time of divorce decrees
- c) ethnic backgrounds of college students
- d) political preferences of corporate executives

Ans: b

2.40 In the following frequency distribution,

LENGTH OF MARRIAGE AT DIVORCE

<u>Length(years)</u>	<u>Percent</u>	<u>Cum Percent</u>
30 or more	2	100
25-29	2	98
20-24	5	96
15-19	8	91
10-14	11	83
5- 9	22	72
0- 4	<u>50</u>	50
	100%	

SOURCE: *Marriage and Divorce: 1988*. Vital Statistics of the United States, Vol. III, 1996.

the total number of observations is

- a) 100
- b) 1,000
- c) unknown.
- d) unknown, but probably quite large, given the source.

Ans: d

2.41 (NOTE: This question requires display of frequency distribution in 2.40.)

This distribution isn't quite as informative as it could be because

- a) only relative frequencies are used.
- b) too many intervals are used.
- c) percents are carried only one digit to the right of the decimal point.
- d) observations create a peculiar lopsided or unbalanced pattern.

Ans: a

2.42 (NOTE: This question requires display of frequency distribution in 2.40.)

The distribution permits us to conclude that

- a) about one-half of all divorced couples are married for fewer than 5 years.
- b) a substantial majority of all divorced couples are married for fewer than 20 years.
- c) a very small minority of all divorced couples are married for 30 or more years.
- d) all of the above

Ans: d

2.43 (NOTE: This question requires display of frequency distribution in 2.40.)

If you could obtain more detailed information about only *one* class, it probably should be

- a) *30 or more*, because this covers the largest number of possible values.
- b) *30 or more*, because the larger values always are more informative.
- c) *0-4*, because this class contains the largest relative frequency.
- d) *0-4*, because the smaller values always are more informative.

Ans: c

2.44 (NOTE: This question requires display of frequency distribution in 2.40.)

The data pattern for this distribution best can be described as

- a) balanced.
- b) lopsided.
- c) either balanced or lopsided, depending on your point of view.
- d) undecipherable because of the absence of regular frequencies.

Ans: b

2.45 Given the following frequency distribution,

SUICIDES BY METHOD, 1992

(in hundreds)

<u>METHOD</u>	<u>MALES</u>	<u>FEMALES</u>
Firearms	163	24
Poison	32	22
Hanging	37	9
Other	<u>15</u>	<u>6</u>
	247	61

SOURCE: *Statistical Abstracts of the U.S.*, 1995.

the number of males who commit suicide with poison equals (BE CAREFUL!)

- a) 32
- b) 320
- c) 3,200
- d) 32,000

Ans: c

2.46 (NOTE: This question requires display of frequency distribution in 2.45.)

Comparison of these two distributions would be facilitated by

- a) converting frequencies to relative frequencies.
- b) expressing the frequencies more exactly.
- c) expanding the *Other* category.
- d) ignoring the *Other* category.

Ans: a

2.47 (NOTE: This question requires display of frequency distribution in 2.45.)

Generally speaking, inspection of the above table suggests that

- a) males and females are about equally likely to commit suicide.
- b) females are about equally likely to use any of the four methods.
- c) males are much more likely to use firearms than females.
- d) males are much more likely to use poison than females.

Ans: c

2.48 Which of the following *is not* a property of histograms?

- a) Units along the vertical axis reflect frequency.
- b) Units along the horizontal axis reflect class intervals of frequency distributions.
- c) Heights of bars reflect the frequencies for the various class intervals.
- d) The intersection of the two axes defines the lower boundary of the bottom class interval.

Ans: d

2.49 Adjacent bars in histograms share common boundaries in order to

- a) produce a more compact graph.
- b) present the data without artificial disruptions.
- c) minimize the effort involved in constructing the graph.
- d) comply with a widely adopted convention.

Ans: b

2.50 When constructing a histogram for the ages of college students, it would be desirable to

- a) locate age -- a type of frequency -- along the vertical axis.
- b) construct only one frequency bar for all students over the age of 40.
- c) break the horizontal scale (with wiggly lines) between 0 and the bottom class interval.
- d) clearly show along the horizontal axis the upper and lower boundaries of every class interval.

Ans: c

2.51 A frequency polygon is

- a) a type of bar graph.
- b) a line graph.
- c) an appropriate graph for qualitative data.
- d) especially appropriate for small batches of data.

Ans: b

2.52 Frequency polygons are particularly useful when

- a) data are qualitative.
- b) data are ungrouped.
- c) two or more frequency distributions are to be displayed on the same graph.
- d) the original frequency distribution is to be portrayed with complete accuracy.

Ans: c

2.53 Histograms and frequency polygons may be viewed as the graphic counterpart of

- a) frequency distributions for quantitative data.
- b) bar graphs.
- c) quantitative data.
- d) class intervals.

Ans: a

- 2.54 A stem and leaf display is ideal for summarizing distributions when you want to
- produce an idealized distribution.
 - compare quantitative and qualitative data.
 - emphasize the orderliness of data.
 - preserve the identities of the individual scores.

Ans: d

- 2.55 A stem and leaf display is a device for sorting data on the basis of
- leading and trailing digits.
 - class intervals.
 - relative frequency.
 - degree of precision.

Ans: a

- 2.56 In a stem and leaf display for annual incomes, an annual income of \$8,234 might be displayed as
- a stem of 8 (thousands) and a leaf of 233.
 - a stem of 4 (units) and a leaf of 823.
 - a stem of 8234 and no leaf.
 - none of the above

Ans: a

- 2.57 An important characteristic of histograms, frequency polygons, and stem and leaf displays is
- size.
 - total area.
 - relative area.
 - shape.

Ans: d

- 2.58 Frequency distributions that approximate a bell-shaped silhouette are described as
- bimodal.
 - normal.
 - positively skewed.
 - negatively skewed.

Ans: b

- 2.59 If the incomes of equal numbers of skilled and unskilled wage earners appear in a single distribution, the resulting frequency distribution probably will be
- bimodal.
 - normal.
 - positively skewed.
 - negatively skewed.

Ans: a

- 2.60 A frequency distribution of standardized IQ scores for equal numbers of male and female grade-school children probably will be
- bimodal.
 - normal.
 - positively skewed.
 - negatively skewed.

Ans: b

- 2.61 A frequency distribution of scores on a very difficult statistics test probably will be
- bimodal.
 - normal.
 - positively skewed.

d) negatively skewed.

Ans: c

2.62 Automobile speeds are clocked along a remote stretch of Interstate 80. If, in fact, motorists are obeying the posted 70 miles per hour speed limit, the frequency distribution of clocked speeds probably will be

- a) bimodal.
- b) normal.
- c) positively skewed.
- d) negatively skewed.

Ans: d

2.63 The term *positively skewed* implies that

a) a *minority* of observations are located in the positive direction relative to the rest of the distribution.

b) a *minority* of observations are positive and the majority of observations are negative.

c) a *majority* of observations are located in the positive direction relative to the rest of the distribution.

d) a *majority* of observations are positive and the minority of observations are negative.

Ans: a

2.64 The feature that distinguishes bar graphs from histograms is the use of

- a) the vertical axis for frequency.
- b) bars.
- c) gaps between bars.
- d) the horizontal axis for class intervals and categories.

Ans: c

2.65 Bar graphs are designed for use with

- a) quantitative data.
- b) qualitative data.
- c) ranked data.
- d) ungrouped data.

Ans: b

2.66 An unscrupulous person could exaggerate frequency differences in histograms or bar graphs by

- a) omitting the lower end of the frequency scale.
- b) increasing the width of taller bars.
- c) making the vertical axis much taller than the width of the horizontal axis.
- d) all of the above

Ans: d

2.67 When constructing graphs, it's considered good practice to make the vertical axis

- a) about as tall as the horizontal axis is wide.
- b) taller than the horizontal axis is wide.
- c) shorter than the horizontal axis is wide.
- d) none of the above

Ans: a

2.68 Although there are many different types of misleading graphs, the book emphasizes those produced by

- a) falsifying data.
- b) withholding data.
- c) unscrupulous graphic techniques.
- d) ignorance.

Ans: c

2.69 When constructing a graph, you must first decide on the type of graph. This decision depends on

- a) the total number of observations.
- b) whether data are quantitative or qualitative.
- c) whether data are grouped or ungrouped.
- d) the impression that you wish to create.

Ans: b

2.70 Wiggly lines can be used to signal scale breaks, and they can appear

- a) along the horizontal axis.
- b) along the vertical axis.
- c) both a and b
- d) neither a nor b

Ans: c

2.71 If you wish to graph the distribution of ethnic backgrounds for a population of alcoholics, use a

- a) histogram.
- b) frequency polygon.
- c) stem and leaf display.
- d) bar graph

Ans: d.

2.72 If you wish to graph the distribution of times spent dreaming during sleep (as inferred from rapid eye movements) for a sample of young adults, use

- a) a histogram or bar graph.
- b) a histogram or frequency polygon.
- c) a bar graph or frequency polygon.
- d) any of the above

Ans: b

2.73 If you wish to use a single bar graph to compare two different frequency distributions -- for instance, one for alcoholics and one for nonalcoholics--it would be essential to

- a) double the frequency scale.
- b) double the width of the horizontal axis.
- c) use some technique, such as shading, to distinguish between the bars for the two distributions.
- d) use some technique, such as the selection of axes that maximizes the differences between the two distributions.

Ans: c

2.74 For a given batch of data, equally competent people might choose

- a) a histogram or frequency polygon.
- b) a histogram or bar graph.
- c) a frequency polygon or bar graph.
- d) any of the above

Ans: a

ANSWERS TO SELECTED QUESTIONS (NOT GIVEN IN TEXTBOOK)

CHAPTER 1

- 1.7** (a) inferential statistics (f) inferential statistics
 (b) descriptive statistics (g) descriptive statistics
 (c) inferential statistics (h) inferential statistics
 (d) descriptive statistics (i) inferential statistics
 (e) inferential statistics
- 1.8** (a) experiment; cages with either one, several, or many other chimps
 (b) observational study
 (c) observational study
 (d) experiment; groups of 2, 3, or 4 students
 (e) observational study
 (f) experiment; test-taking or control workshop; allowing students to select type of workshop might confound results
 (g) observational study
 (h) observational study

CHAPTER 2

2.14 Residences

	<i>f</i>
9	1
8	0
7	1
6	1
5	1
4	5
3	8
2	6
<u>1</u>	<u>2</u>
Total	25

- 2.15 (a) positively skewed**
 (b)

<u>FRIENDS</u>	<u>Relative <i>f</i></u>
400 – above	.01
350 – 399	.03
300 – 349	.06
250 – 299	.09
200 – 249	.12
150 – 199	.25
100 – 149	.14
50 – 99	.15
<u>0 – 49</u>	<u>.18</u>
Total	1.06

- (c) 96.5 approximate percentile rank (Calculated from freq. table in book.)
 (d) Individual observations not given

- 2.16** (a) The unit of measurement equals one-hundredth of a point (.01).
 (b) For nonmediators,

$$\text{class width} = \frac{3.79 - 1.90}{10} = \frac{1.89}{10} = .189.$$

Round off .189 to .20.

GPA	Frequency	
	Nonmediators	Mediators
3.60 – 3.79	2	3
3.40 – 3.59	2	8
3.20 – 3.39	3	5
3.00 – 3.19	4	3
2.80 – 2.99	5	2
2.60 – 2.79	6	2
2.40 – 2.59	2	2
2.20 – 2.39	1	2
2.00 – 2.19	1	0
1.80 – 1.99	<u>1</u>	<u>0</u>
Totals	27	27

(c) Mediators tend to have higher GPAs than nonmediators.

- 2.18** (a) 68,300
(b)

Major Field	Percent	
	Males	Females
Business	32	23
Social studies	16	12
Education	4	13
Health sciences	3	12
Psychology	4	10
Engineering	13	2
Life sciences	5	6
Fine Arts	6	8
Communications	6	7
Computer sciences	7	1
English	3	6
Totals	<u>99%</u>	<u>100%</u>

Among male graduates, there are relatively more degrees particularly in engineering and business, whereas among female graduates, there are relatively more degrees particularly in education and health sciences.

- (c) No, because there is no obvious order associated with the major fields of study.
(d)

Bachelor's Degrees Conferred by Study Field and Gender (2005-2006)

