

MN 576 NEW GENERATION FINAL EXAM FOR WEEK 10

Question 1

3 / 3 points

Elevated inhibin-A is noted when a pregnant woman is at increased risk of having an infant with:

Question options:

1)

Down syndrome.

2)

Edwards syndrome.

3)

open neural tube defect.

4)

hemolytic anemia.

Question 2

3 / 3 points

During well-women visits for 21- to 29-year-old sexually-active women who report more than 1 sex partner within the past 6 months, all of the following are appropriate screening tests except:

Question options:

1)

Pap smear.

2)

HPV testing.

3)

Pelvic examination.

4)

STI screening.

Question 3

3 / 3 points

Components of the antenatal screening test known as the "quad screen" include all of the following except:

Question options:

1)

AFP.

2)

hCG.

3)

unconjugated estriol.

4)

progesterone.

Question 4

1 / 3 points

Anxiety in response to a challenging life event is a natural response by the body to:

Question options:

1)

help a person focus on the issue at hand.

2)

diminish the fight-or-flight response.

3)

impair decision-making under duress.

4)

provide transient improvement in physical capabilities.

Question 5

3 / 3 points

Women who experienced gestational diabetes are at higher risk of developing Type 2 Diabetes later in life.

Question options:

1) True

2) False

Question 6

0 / 3 points

Malignant thyroid nodules are more common in women than men.

Question options:

1) True

2) False

Question 7

3 / 3 points

When considering characteristics of the domestic violence perpetrator. One of the best predictors of a subsequent homicide of victims of domestic violence is which of the following?

Question options:

1)

history of perpetrator striking victim on the face with an open hand

2)

history of perpetrator attempting to strangle the victim

3)

perpetrator's access to kitchen knives.

4)

alcohol abuse history in the victim

Question 8

3 / 3 points

Treatment of vulvovaginitis caused by *Candida albicans* includes:

Question options:

1)

metronidazole gel.

2)

clotrimazole cream.

3)

hydrocortisone ointment.

4)

clindamycin cream.

Question 9

3 / 3 points

HDL levels are a better predictor of CAD in women than LDL.

Question options:

1) True

2) False

Question 10

3 / 3 points

Diagnostic criteria for generalized anxiety disorder include all of the following except:

Question options:

1)

difficulty concentrating.

2)

consistent early morning wakening.

3)

apprehension.

4)

irritability.

Question 11

3 / 3 points

Concerning the use of alcohol during pregnancy, which of the following statements is most accurate?

Question options:

1)

Although potentially problematic, maternal alcohol intake does not increase the risk of miscarriage.

2)

Risk to the fetus from alcohol exposure is greatest in the third trimester.

3)

No level or time of exposure is considered to be safe.

4)

Risk of fetal alcohol syndrome is present only if alcohol exposure has occurred throughout the pregnancy.

Question 12

3 / 3 points

Which of the following is most consistent with the diagnosis of generalized anxiety disorder?

Question options:

1)

gastrointestinal upset

2)

difficulty initiating sleep

3)

diminished cognitive ability

4)

consistent early morning wakening

Question 13

3 / 3 points

For a woman with bothersome hot flashes who cannot take HT, alternative options with demonstrated efficacy and limited adverse effects include the use of all of the following except:

Question options:

1)

venlafaxine.

2)

sertraline.

3)

gabapentin.

4)