

NCLEX PN 2023 Study Guide Exam Questions with Verified solutions Latest Update.

Adverse effect of Erythropoietin? (Used to treat anemia) - **Correct**

Answer: Hypertension

Adverse effects of INH (Isoniazid) - **Correct Answer:** • Hepatotoxicity (skin jaundice, vomiting, dark urine)

- Peripheral neuropathy (numbness, tingling.)

DO NOT DRINK WITH ALCOHOL

TREATS TB!!!

Antiplatelet agents - **Correct Answer:** Substance that interferes with the action of platelets. Prolongs bleeding time. Used to prevent heart attacks and strokes.

ANTIPLATELET AGENTS' NAMES - **Correct Answer:** Clopidogrel, Ticagrelor, Prasugrel, Aspirin.

Antiplatelet agents should not be taken by? - **Correct Answer:** Clients with bleeding peptic ulcers, active bleeding, or intracranial hemorrhage.

Drinks to avoid while on warfarin therapy. - **Correct Answer:** Green tea, grapefruit juice, and cranberry juice.

Factors that increase risk of falls - **Correct Answer:** • Positive orthostatic vital signs

- Carbidopa/Levodopa (Parkinson's meds)
- > 65yr old
- Using assistive devices
- Osteoarthritis of knees

Foods you should avoid excess amounts/ inconsistent intake of with warfarin therapy. - **Correct Answer:** Vitamin K rich foods (leafy green vegetables - broccoli, kale, asparagus, & spinach)

How far should you insert enema tube into rectum? - **Correct Answer:** 3-4in

Major adverse effects of statin medications (Atorvastatin) - **Correct Answer:** • Muscle cramps
• Liver injuries

Major side effects of Metformin - **Correct Answer:** decreased in vitamin b12 levels

lactic acidosis

(other non-major N/V, diarrhea, epigastric pain)

meds that might cause orthostatic hypotension - **Correct Answer:** Antihypertensive, diuretics, antipsychotics, antidepressants, and narcotics.

Normal HR - **Correct Answer:** 60-100 bpm

Normal platelet range - **Correct Answer:** 150,000-400,000

Proper cane length calls for the cane to be level with the user's greater trochanter and for the user's elbow to be bent at a: - **Correct Answer:** 20-30degree angle + hold the cane 6-10in in front of them.

Risk factors for MRSA - **Correct Answer:** • Recent or prolonged hospitalization

- Residence in a long term care facility
- Invasive lines (Hemodialysis catheter)
- Sharing needles, razors, or sports equipment
- Advanced age
- Antibiotic use

Some names of tumor necrosis factor (TNF) inhibitors - **Correct Answer:** (Infliximab, adalimumab, etanercept)

Therapeutic Lithium level - **Correct Answer:** 0.6-1.2

Therapeutic response of Levothyroxine - **Correct Answer:** • Increase energy levels

- HR WNL

-

Tumor necrosis factors (TNF) - **Correct Answer:** Suppresses the inflammatory response in autoimmune diseases such as RA, Chron disease, and psoriasis. Due to the immunosuppressive action of these drugs clients are at increased risk of infection.

What type of clients should not take TNF's? - **Correct Answer:** Clients with current, recent, or chronic infections.

Which immediate acting meds to give for an acute asthma attack? - **Correct Answer:** Albuterol and Ipratropium

Which side should you hold the cane on? - **Correct Answer:** stronger side