

sabado

- 1. A home health nurse is providing teaching about post seizure management to the parents of a school age child who has epilepsy and experiences tonic-clonic seizures. Which of the following instructions should the nurse include in the teaching? (Select all that apply.)**

"Hyperextend your child's head for 5 minutes following a seizure."

"Immediately following a seizure, give your child 6 ounces of water."

"Following a seizure, record the length and characteristics of your child's seizure."

"Administer rectal diazepam to your child following a seizure."

"Call for emergency medical services if the size of your child's pupils are unequal after a seizure."

- 2. A nurse in a pediatric clinic is discussing the pathophysiology of Reye syndrome with a newly licensed nurse. Which of the following statements by the newly licensed nurse indicate an understanding of the disorder?**

"Reye syndrome causes fatty changes in the liver."

"Reye syndrome leads to venous thrombus formation."

"Reye syndrome is associated with misuse of acetaminophen."

"Reye syndrome is linked to decreased serum ammonia levels"

- 3. A nurse is teaching the parents of a toddler about strategies to manage temper tantrums. Which of the following instructions should the nurse include in the teaching?**

"Honor the child's request if she holds her breath."

"Establish a structured daily routine for the child."

"Place the child in her room alone until the temper tantrum ends."

"Comfort the child during the temper tantrum."

- 4. A nurse is completing an assessment following suctioning of a child who has a tracheostomy. Which of the following findings should the nurse identify as an indication that the procedure has been effective?**

Increased respiratory rate

Decreased oxygen saturation

Clear breath sounds

Increased oral secretions

- 5. A nurse is developing a plan of care for a 4-year-old child who has hemophilia and is experiencing acute hemarthrosis. Which of the following interventions should the nurse include in the plan?**

Have the child perform passive range-of-motion exercises.

Administer aspirin as needed for pain.

Place ice packs on the affected joints.

Position the lower extremities below the level of the heart.

- 6. A nurse in the emergency department is caring for a preschooler who has epiglottitis. Which of the following actions should the nurse take?**

Place the child in a left lateral position.

Obtain a specimen from the child's throat for a culture.

Inspect the child's throat with a padded tongue depressor.

Initiate droplet precautions for the child.

- 7. A nurse is providing teaching to the parent of a child about home care following a cardiac catheterization. Which of the following statements should the nurse include?**

"You should give your child aspirin if they report pain at the site."

"You can remove the pressure dressing 8 hours after the procedure."

"Your child can soak in the bathtub 24 hours after the procedure."

"Your child should avoid strenuous physical activities for several days."

- 8. A nurse is admitting a child who has pertussis. Which of the following isolation precautions should the nurse initiate?**

Droplet

Contact

Airborne

Protective environment

**9. A nurse on a pediatric unit is receiving change-of-shift report for a group of clients. Which of the following clients should the nurse see first?**

A child who is postoperative following a tonsillectomy and reports moderate throat pain

A child who had a cardiac catheterization using the femoral artery and has blanching of the toes

A child who has bacterial pneumonia and is due for their initial dose of IV antibiotics

A child who has juvenile idiopathic arthritis and needs assistance with application of prescribed splints

**10. A nurse in a pediatric clinic is performing a history and physical for a toddler who is scheduled to receive a measles, mumps, and rubella (MMR) immunization. Which of the following findings indicate that the immunization should be withheld?**

Temperature of 38° C (100.4° F)

Family history of sudden infant death syndrome (SIDS)

Taking an antihistamine for seasonal allergies

Receiving prednisone for nephrotic syndrome

**11. A nurse is instilling optic drops into an 18-month-old child's ears. Which of the following methods should the nurse use?**

Pull the pinna down and back.

Insert the dropper into the ear canal.

Administer the ear drops at 5.5° C (42° F).

Massage the area behind the ear.

**12. A nurse is providing teaching to an adolescent who has a new prescription for cefazolin. For which of the following adverse effects should the nurse instruct the adolescent to monitor and report to the provider?**

Dry mouth

Constipation

Back pain

## Urticaria

**13. A nurse is providing discharge teaching to the parents of a child who has sickle cell anemia and was hospitalized after experiencing a vasoocclusive crisis. Which of the following instructions should the nurse include in the teaching?**

"Avoid immunizing your child with the meningococcal vaccine."

"Restrict the time your child spends playing outdoors."

**"Increase your child's intake of oral fluids."**

"Provide your child with a high-protein diet."

**14. A nurse is providing teaching to the parent of a school-age child who has pediculosis capitis. Which of the following instructions should the nurse include in the teaching?**

"Seal no washable items in a plastic bag for 2 days."

**"Soak hair brushes in boiling water for 10 minutes."**

"Apply permethrin 1 percent cream rinse every day for 5 days."

"After washing bed linens, place them in a dryer on a cool setting for 30 minutes."

**15. A nurse is admitting a child who has bacterial meningitis. Which of the following actions should the nurse take first?**

**Initiate antibiotic therapy for the child.**

Minimize the child's environmental stimuli.

Place the child in a side-lying position.

Administer pain medication to the child

**16. A nurse is assessing a 12-month-old infant who is immediately postoperative following hernia repair surgery. Which of the following pain measurement tools should the nurse use to determine if the infant is experiencing pain?**

FACES

COMFORT

CRIS

**FLACC**

**17. A nurse is caring for a school-age child who has metastatic osteosarcoma. The child asks the nurse, "Am I going to die?" Which of the following responses should the nurse make?**

"What is your pain level right now?"

"Your doctor will be able to answer your questions tomorrow."

"It sounds like you are worried. Tell me what you have been told."

"It's natural to worry about death, but you should focus your energy on getting better."

**18. A nurse is caring for a toddler who drinks 946 mL (32 oz) of whole milk per day and has a poor appetite. The nurse should identify the toddler as being at risk for which of the following conditions?**

Iron deficiency anemia

Vitamin A toxicity

Impaired carbohydrate metabolism

Lactose intolerance

**19. A nurse is providing teaching to the parent of a child who has a new prescription for methylphenidate. Which of the following instructions should the nurse include?**

"Monitor your child for excessive sleepiness."

"Administer the medication with a caffeinated beverage."

"Administer the second dose of the medication at lunch time."

"Monitor your child for weight gain."

**20. A nurse is planning care for an infant who has congenital talipes equinovarus. Which of the following referrals should the nurse expect the provider to make?**

Ophthalmologist

Nutrition counselor

Speech therapist

Orthopedic specialist

**21. A nurse is caring for a child who has acute glomerulonephritis. Which of the following actions should the nurse take?**