

Nurs 6635 || Psychopathology exam || RATED A|| LATEST UPDATE 2023/24||ALL YOU NEED.

Question 3

What is the recommended duration of treatment with psychopharmacology of panic disorder once treatment becomes effective?

- 4-6 weeks
- 1-2 months
- 3-7 months
- 8-12 months

Question 4

Select age after which encopresis may be correctly diagnosed.

- 2
- 3
- 4
- 6

Question 5

Select the antidepressant drug that is most likely to cause death when taken in an overdose.

- Fluoxetine
- Mirtazapine
- Imipramine
- Trazodone

Question 6

Select the disorder in which inflicting injury to self or others is common.

- Pain
- Conversion

- Somatic symptom
- Factitious

Question 7

- A. 0.1-0.5
- B. 0.6-1.2
- C. 1.4-1.9
- D. 2.0-2.6

Question 9

Select the antidepressant drug that inhibits reuptake of both norepinephrine and serotonin (5-HT).

- Fluoxetine
- Desipramine
- Venlafaxine
- Bupropion

Question 10

Select the psychiatric disorder that occurs most commonly in patients hospitalized for medical problems and surgery.

- Depression
- Adjustment
- Bipolar

Short
D.
Anxiety

Question 11

Select the short rating scale used to assess change in psychiatric inpatients.

- PANSS

- BPRS
- HAM-D
- CAPS

Question 12

- Opioid analgesics
- TCA and SSRI antidepressants
- Beta-adrenergic blockers
- Mood stabilizers

Question 13

Select the medical condition that must be excluded as a cause of somatic symptom disorder or hypochondriasis.

- Angina pectoris
- Asthma
- Diabetes
- Multiple sclerosis

Question 14

Select the most commonly occurring eating disorder.

- Anorexia nervosa
- Bulimia nervosa
- Purging
- Binge

Question 15

Select the age stage of life in which identity crisis typically emerges.

- Childhood
- Adolescence

- Middle adult
- Elderly adult

Question 16

Select the **two** rating scales used specifically for schizophrenia.

- PANSS
- MMSE
- SAPS
- BPRS

Question 17

Select the drug that has been effective in treating social anxiety in children.

- Bupropion
- Carbamazepine
- Amitriptyline
- Fluoxetine

Question 18

Select the drug used to treat bipolar disorder that may cause hypothyroidism.

- Lithium
- Risperidone
- Lamotrigine
- Valproate

Question 19

Select the neurotransmitter that is most genetically involved in childhood and adolescent depression.

- Norepinephrine
- Serotonin
- Dopamine

- Gamma amino butyric acid (GABA)

Question 20

Select the most likely stressor to cause an adjustment disorder.

- Natural disaster
- Emergency surgery
- Vehicle accident
- Substance abuse

Question 21

Select the cognitive function category in a mental status examination that is represented by the question, "Starting at 3 A.M., count back the time 2 hours at a time."

- Orientation
- Concentration
- Calculation
- Alertness

Question 21

Select the cognitive function category in a mental status examination that is represented by the question, "Starting at 3 A.M., count back the time 2 hours at a time."

- Orientation
- Concentration
- Calculation
- Alertness

Question 22

Select the memory category retained in transient global amnesia that is absent in dissociative amnesia.

- Family members
- Sequential events